

表 3524 PTH値(pg/mL) HDL-C濃度(mg/dL)別 (施設血液透析患者 透析回数週3回) PTH測定法:intact-PTH

HDL-C濃度	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
20未満	75	72	64	48	46	39	31	36	28	46	2	47	21	8	7	2	1		3	3	579	19	598	128.27	130.04
(%)	(13.0)	(12.4)	(11.1)	(8.3)	(7.9)	(6.7)	(5.4)	(6.2)	(4.8)	(7.9)	(0.3)	(8.1)	(3.6)	(1.4)	(1.2)	(0.3)	(0.2)		(0.5)	(0.5)	(100.0)				
20～	637	840	865	920	923	906	830	817	744	1,605	23	932	489	248	162	49	69	78	119	55	11,311	173	11,484	162.68	141.13
(%)	(5.6)	(7.4)	(7.6)	(8.1)	(8.2)	(8.0)	(7.3)	(7.2)	(6.6)	(14.2)	(0.2)	(8.2)	(4.3)	(2.2)	(1.4)	(0.4)	(0.6)	(0.7)	(1.1)	(0.5)	(100.0)				
30～	1,592	2,367	2,424	2,584	2,593	2,691	2,544	2,271	2,055	4,716	58	2,687	1,531	846	473	138	206	229	309	255	32,569	482	33,051	168.23	151.95
(%)	(4.9)	(7.3)	(7.4)	(7.9)	(8.0)	(8.3)	(7.8)	(7.0)	(6.3)	(14.5)	(0.2)	(8.3)	(4.7)	(2.6)	(1.5)	(0.4)	(0.6)	(0.7)	(0.9)	(0.8)	(100.0)				
40～	1,760	2,467	2,743	2,872	2,921	2,887	2,785	2,605	2,280	5,144	71	2,989	1,696	984	562	133	231	235	380	309	36,054	426	36,480	170.54	161.75
(%)	(4.9)	(6.8)	(7.6)	(8.0)	(8.1)	(8.0)	(7.7)	(7.2)	(6.3)	(14.3)	(0.2)	(8.3)	(4.7)	(2.7)	(1.6)	(0.4)	(0.6)	(0.7)	(1.1)	(0.9)	(100.0)				
50～	1,169	1,794	1,935	2,119	2,143	1,959	1,971	1,803	1,678	3,672	37	2,118	1,222	748	425	110	155	155	307	244	25,764	295	26,059	173.57	171.86
(%)	(4.5)	(7.0)	(7.5)	(8.2)	(8.3)	(7.6)	(7.7)	(7.0)	(6.5)	(14.3)	(0.1)	(8.2)	(4.7)	(2.9)	(1.6)	(0.4)	(0.6)	(0.6)	(1.2)	(0.9)	(100.0)				
60～	630	973	1,049	1,245	1,210	1,202	1,111	1,032	985	2,153	22	1,298	774	408	262	35	91	106	161	126	14,873	192	15,065	174.14	162.22
(%)	(4.2)	(6.5)	(7.1)	(8.4)	(8.1)	(8.1)	(7.5)	(6.9)	(6.6)	(14.5)	(0.1)	(8.7)	(5.2)	(2.7)	(1.8)	(0.2)	(0.6)	(0.7)	(1.1)	(0.8)	(100.0)				
70～	313	478	549	548	593	593	556	527	481	1,100	15	599	348	165	112	29	43	66	86	65	7,266	75	7,341	173.37	162.01
(%)	(4.3)	(6.6)	(7.6)	(7.5)	(8.2)	(8.2)	(7.7)	(7.3)	(6.6)	(15.1)	(0.2)	(8.2)	(4.8)	(2.3)	(1.5)	(0.4)	(0.6)	(0.9)	(1.2)	(0.9)	(100.0)				
80～	151	239	260	296	240	292	254	236	213	479	6	260	146	105	50	11	27	26	37	24	3,352	37	3,389	170.09	160.53
(%)	(4.5)	(7.1)	(7.8)	(8.8)	(7.2)	(8.7)	(7.6)	(7.0)	(6.4)	(14.3)	(0.2)	(7.8)	(4.4)	(3.1)	(1.5)	(0.3)	(0.8)	(0.8)	(1.1)	(0.7)	(100.0)				
90～	53	101	111	125	133	123	123	98	102	207	4	144	78	47	24	10	7	14	23	15	1,542	8	1,550	180.43	169.66
(%)	(3.4)	(6.5)	(7.2)	(8.1)	(8.6)	(8.0)	(8.0)	(6.4)	(6.6)	(13.4)	(0.3)	(9.3)	(5.1)	(3.0)	(1.6)	(0.6)	(0.5)	(0.9)	(1.5)	(1.0)	(100.0)				
100～	67	91	103	87	125	108	93	86	81	209	3	95	56	46	22	1	7	12	16	10	1,318	13	1,331	170.40	148.12
(%)	(5.1)	(6.9)	(7.8)	(6.6)	(9.5)	(8.2)	(7.1)	(6.5)	(6.1)	(15.9)	(0.2)	(7.2)	(4.2)	(3.5)	(1.7)	(0.1)	(0.5)	(0.9)	(1.2)	(0.8)	(100.0)				
合計	6,447	9,422	10,103	10,844	10,927	10,800	10,298	9,511	8,647	19,331	241	11,169	6,361	3,605	2,099	518	837	921	1,441	1,106	134,628	1,720	136,348	170.37	159.74
(%)	(4.8)	(7.0)	(7.5)	(8.1)	(8.1)	(8.0)	(7.6)	(7.1)	(6.4)	(14.4)	(0.2)	(8.3)	(4.7)	(2.7)	(1.6)	(0.4)	(0.6)	(0.7)	(1.1)	(0.8)	(100.0)				
記載なし	3,065	4,307	4,422	4,670	4,744	4,471	4,288	3,992	3,460	8,097	97	4,429	2,631	1,519	925	218	352	389	691	516	57,283	2,955	60,238	169.03	163.82
(%)	(5.4)	(7.5)	(7.7)	(8.2)	(8.3)	(7.8)	(7.5)	(7.0)	(6.0)	(14.1)	(0.2)	(7.7)	(4.6)	(2.7)	(1.6)	(0.4)	(0.6)	(0.7)	(1.2)	(0.9)	(100.0)				
総計	9,512	13,729	14,525	15,514	15,671	15,271	14,586	13,503	12,107	27,428	338	15,598	8,992	5,124	3,024	736	1,189	1,310	2,132	1,622	191,911	4,675	196,586	169.97	160.97
(%)	(5.0)	(7.2)	(7.6)	(8.1)	(8.2)	(8.0)	(7.6)	(7.0)	(6.3)	(14.3)	(0.2)	(8.1)	(4.7)	(2.7)	(1.6)	(0.4)	(0.6)	(0.7)	(1.1)	(0.8)	(100.0)				
平均	47.00	47.72	48.16	48.27	48.40	48.32	48.30	48.17	48.48	48.42	48.13	48.33	48.68	49.09	48.99	46.75	48.27	49.27	49.61	49.53	48.29	46.08	48.26		
標準偏差	16.75	16.63	16.81	16.44	16.60	16.67	16.43	16.32	16.55	16.63	17.52	16.45	16.44	16.67	16.53	15.57	16.38	17.03	17.26	15.68	16.57	16.04	16.57		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査