

表 3526 PTH値(pg/mL) 透析前収縮期血圧(mmHg)別（施設血液透析患者 透析回数週3回）PTH測定法:intact-PTH

透析前収縮期血圧	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
60未満	1	2		2	3	3	1	3	3	4		2	1	1	1						27	2	29	166.15	107.58
(%)	(3.7)	(7.4)		(7.4)	(11.1)	(11.1)	(3.7)	(11.1)	(11.1)	(14.8)		(7.4)	(3.7)	(3.7)	(3.7)					(100.0)					
60～	16	25	23	15	15	19	7	17	10	32		18	11	3	4	1		3	5	2	226	9	235	164.93	161.57
(%)	(7.1)	(11.1)	(10.2)	(6.6)	(6.6)	(8.4)	(3.1)	(7.5)	(4.4)	(14.2)		(8.0)	(4.9)	(1.3)	(1.8)	(0.4)		(1.3)	(2.2)	(0.9)	(100.0)				
80～	143	146	168	156	191	156	147	132	114	252	1	139	99	50	22	12	20	7	32	29	2,016	48	2,064	172.04	198.94
(%)	(7.1)	(7.2)	(8.3)	(7.7)	(9.5)	(7.7)	(7.3)	(6.5)	(5.7)	(12.5)	(0.0)	(6.9)	(4.9)	(2.5)	(1.1)	(0.6)	(1.0)	(0.3)	(1.6)	(1.4)	(100.0)				
100～	643	861	907	968	923	859	792	771	723	1,487	12	872	449	302	181	47	80	82	140	125	11,224	303	11,527	169.33	169.70
(%)	(5.7)	(7.7)	(8.1)	(8.6)	(8.2)	(7.7)	(7.1)	(6.9)	(6.4)	(13.2)	(0.1)	(7.8)	(4.0)	(2.7)	(1.6)	(0.4)	(0.7)	(0.7)	(1.2)	(1.1)	(100.0)				
120～	1,753	2,547	2,670	2,794	2,893	2,861	2,770	2,503	2,279	5,198	70	2,907	1,678	931	564	120	228	248	392	308	35,714	781	36,495	170.75	165.33
(%)	(4.9)	(7.1)	(7.5)	(7.8)	(8.1)	(8.0)	(7.8)	(7.0)	(6.4)	(14.6)	(0.2)	(8.1)	(4.7)	(2.6)	(1.6)	(0.3)	(0.6)	(0.7)	(1.1)	(0.9)	(100.0)				
140～	2,635	3,870	4,135	4,516	4,537	4,596	4,303	3,979	3,572	8,058	92	4,577	2,576	1,414	842	203	323	384	549	388	55,549	1,050	56,599	167.50	150.27
(%)	(4.7)	(7.0)	(7.4)	(8.1)	(8.2)	(8.3)	(7.7)	(7.2)	(6.4)	(14.5)	(0.2)	(8.2)	(4.6)	(2.5)	(1.5)	(0.4)	(0.6)	(0.7)	(1.0)	(0.7)	(100.0)				
160～	1,958	2,890	3,110	3,304	3,339	3,228	3,145	2,853	2,587	5,845	80	3,321	1,980	1,118	624	163	228	254	432	329	40,788	716	41,504	169.59	157.64
(%)	(4.8)	(7.1)	(7.6)	(8.1)	(8.2)	(7.9)	(7.7)	(7.0)	(6.3)	(14.3)	(0.2)	(8.1)	(4.9)	(2.7)	(1.5)	(0.4)	(0.6)	(0.6)	(1.1)	(0.8)	(100.0)				
180～	785	1,113	1,114	1,276	1,293	1,239	1,138	1,088	953	2,210	27	1,302	743	440	271	67	94	128	174	129	15,584	288	15,872	171.87	158.81
(%)	(5.0)	(7.1)	(7.1)	(8.2)	(8.3)	(8.0)	(7.3)	(7.0)	(6.1)	(14.2)	(0.2)	(8.4)	(4.8)	(2.8)	(1.7)	(0.4)	(0.6)	(0.8)	(1.1)	(0.8)	(100.0)				
200～	201	304	321	282	327	300	286	244	264	526	12	319	201	117	64	23	36	32	60	31	3,950	63	4,013	175.35	175.91
(%)	(5.1)	(7.7)	(8.1)	(7.1)	(8.3)	(7.6)	(7.2)	(6.2)	(6.7)	(13.3)	(0.3)	(8.1)	(5.1)	(3.0)	(1.6)	(0.6)	(0.9)	(0.8)	(1.5)	(0.8)	(100.0)				
220～	33	70	59	51	61	65	53	53	47	99	1	69	27	35	13	6	11	4	12	8	777	16	793	181.44	185.78
(%)	(4.2)	(9.0)	(7.6)	(6.6)	(7.9)	(8.4)	(6.8)	(6.8)	(6.0)	(12.7)	(0.1)	(8.9)	(3.5)	(4.5)	(1.7)	(0.8)	(1.4)	(0.5)	(1.5)	(1.0)	(100.0)				
合計	8,168	11,828	12,507	13,364	13,582	13,326	12,642	11,643	10,552	23,711	295	13,526	7,765	4,411	2,586	642	1,020	1,142	1,796	1,349	165,855	3,276	169,131	169.55	159.05
(%)	(4.9)	(7.1)	(7.5)	(8.1)	(8.2)	(8.0)	(7.6)	(7.0)	(6.4)	(14.3)	(0.2)	(8.2)	(4.7)	(2.7)	(1.6)	(0.4)	(0.6)	(0.7)	(1.1)	(0.8)	(100.0)				
記載なし	1,344	1,901	2,018	2,150	2,089	1,945	1,944	1,860	1,555	3,717	43	2,072	1,227	713	438	94	169	168	336	273	26,056	1,399	27,455	172.63	172.66
(%)	(5.2)	(7.3)	(7.7)	(8.3)	(8.0)	(7.5)	(7.5)	(7.1)	(6.0)	(14.3)	(0.2)	(8.0)	(4.7)	(2.7)	(1.7)	(0.4)	(0.6)	(0.6)	(1.3)	(1.0)	(100.0)				
総計	9,512	13,729	14,525	15,514	15,671	15,271	14,586	13,503	12,107	27,428	338	15,598	8,992	5,124	3,024	736	1,189	1,310	2,132	1,622	191,911	4,675	196,586	169.97	160.97
(%)	(5.0)	(7.2)	(7.6)	(8.1)	(8.2)	(8.0)	(7.6)	(7.0)	(6.3)	(14.3)	(0.2)	(8.1)	(4.7)	(2.7)	(1.6)	(0.4)	(0.6)	(0.7)	(1.1)	(0.8)	(100.0)				
平均	151.03	151.64	151.52	151.60	151.80	151.78	151.76	151.65	151.62	151.84	154.19	152.20	152.46	152.77	152.20	153.44	151.51	152.41	151.73	149.87	151.78	149.01	151.73		
標準偏差	24.84	24.52	24.20	23.83	24.13	23.75	23.46	23.63	23.81	23.55	23.52	23.75	23.82	24.60	24.04	25.92	26.08	24.71	25.92	25.84	23.98	24.66	24.00		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査