

表 3532 PTH値(pg/mL) 原疾患別（血液透析濾過患者 透析回数週3回）PTH測定法intact-PTH

原疾患	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	714	796	856	923	1,003	995	971	887	783	1,926	33	1,124	650	440	261	73	96	112	200	187	13,030	235	13,265	187.70	193.25
(%)	(5.5)	(6.1)	(6.6)	(7.1)	(7.7)	(7.6)	(7.5)	(6.8)	(6.0)	(14.8)	(0.3)	(8.6)	(5.0)	(3.4)	(2.0)	(0.6)	(0.7)	(0.9)	(1.5)	(1.4)	(100.0)				
慢性腎盂腎炎	23	20	29	17	26	27	32	24	15	56	1	30	21	11	9	2	3	2	8	9	365	5	370	202.65	213.92
(%)	(6.3)	(5.5)	(7.9)	(4.7)	(7.1)	(7.4)	(8.8)	(6.6)	(4.1)	(15.3)	(0.3)	(8.2)	(5.8)	(3.0)	(2.5)	(0.5)	(0.8)	(0.5)	(2.2)	(2.5)	(100.0)				
急速進行性糸球体腎炎	14	20	17	26	19	17	10	11	23	33	1	17	12	16	5		4	2			247	6	253	166.24	125.70
(%)	(5.7)	(8.1)	(6.9)	(10.5)	(7.7)	(6.9)	(4.0)	(4.5)	(9.3)	(13.4)	(0.4)	(6.9)	(4.9)	(6.5)	(2.0)		(1.6)	(0.8)			(100.0)				
妊娠腎/妊娠中毒症	20	15	36	25	24	19	22	18	22	40		20	21	5	7	1	3	2	6	6	312	6	318	185.15	207.21
(%)	(6.4)	(4.8)	(11.5)	(8.0)	(7.7)	(6.1)	(7.1)	(5.8)	(7.1)	(12.8)		(6.4)	(6.7)	(1.6)	(2.2)	(0.3)	(1.0)	(0.6)	(1.9)	(1.9)	(100.0)				
その他の分類不能の腎炎	9	11	7	10	12	17	13	14	7	30		19	6	3	4		1	4	5	2	174	2	176	192.66	167.12
(%)	(5.2)	(6.3)	(4.0)	(5.7)	(6.9)	(9.8)	(7.5)	(8.0)	(4.0)	(17.2)		(10.9)	(3.4)	(1.7)	(2.3)		(0.6)	(2.3)	(2.9)	(1.1)	(100.0)				
多発性嚢胞腎	58	69	80	77	96	95	110	77	84	187	4	109	73	52	20	6	10	9	11	13	1,240	23	1,263	183.00	151.01
(%)	(4.7)	(5.6)	(6.5)	(6.2)	(7.7)	(7.7)	(8.9)	(6.2)	(6.8)	(15.1)	(0.3)	(8.8)	(5.9)	(4.2)	(1.6)	(0.5)	(0.8)	(0.7)	(0.9)	(1.0)	(100.0)				
腎硬化症	96	163	175	193	211	206	172	178	152	374	8	235	134	77	34	7	22	21	26	25	2,509	53	2,562	179.43	171.53
(%)	(3.8)	(6.5)	(7.0)	(7.7)	(8.4)	(8.2)	(6.9)	(7.1)	(6.1)	(14.9)	(0.3)	(9.4)	(5.3)	(3.1)	(1.4)	(0.3)	(0.9)	(0.8)	(1.0)	(1.0)	(100.0)				
悪性高血圧	13	11	22	10	18	31	23	20	11	36		25	19	8	7	1	1	2	4	5	267	6	273	192.64	171.52
(%)	(4.9)	(4.1)	(8.2)	(3.7)	(6.7)	(11.6)	(8.6)	(7.5)	(4.1)	(13.5)		(9.4)	(7.1)	(3.0)	(2.6)	(0.4)	(0.4)	(0.7)	(1.5)	(1.9)	(100.0)				
糖尿病性腎症	508	777	852	927	1,003	949	875	795	746	1,629	22	972	529	295	156	38	63	60	103	53	11,352	297	11,649	163.76	137.08
(%)	(4.5)	(6.8)	(7.5)	(8.2)	(8.8)	(8.4)	(7.7)	(7.0)	(6.6)	(14.3)	(0.2)	(8.6)	(4.7)	(2.6)	(1.4)	(0.3)	(0.6)	(0.5)	(0.9)	(0.5)	(100.0)				
SLE腎炎	15	25	15	21	29	25	18	25	17	47		24	20	13	3		2	4	4	2	309	8	317	184.24	236.91
(%)	(4.9)	(8.1)	(4.9)	(6.8)	(9.4)	(8.1)	(5.8)	(8.1)	(5.5)	(15.2)		(7.8)	(6.5)	(4.2)	(1.0)		(0.6)	(1.3)	(1.3)	(0.6)	(100.0)				
アミロイド腎		4	6	4	4	7	6	4	1	7		2			1	1			1		48	2	50	141.08	110.61
(%)		(8.3)	(12.5)	(8.3)	(8.3)	(14.6)	(12.5)	(8.3)	(2.1)	(14.6)		(4.2)		(2.1)	(2.1)				(2.1)		(100.0)				
痛風腎	10	9	12	9	9	8	4	6	9	20	2	16	6	5	4	1			3	1	134	4	138	181.50	150.90
(%)	(7.5)	(6.7)	(9.0)	(6.7)	(6.7)	(6.0)	(3.0)	(4.5)	(6.7)	(14.9)	(1.5)	(11.9)	(4.5)	(3.7)	(3.0)	(0.7)			(2.2)	(0.7)	(100.0)				
先天性代謝異常による腎不全	2	2	2	4	3	4	1	2	3	4		2	3	3						1	36		36	178.58	170.02
(%)	(5.6)	(5.6)	(5.6)	(11.1)	(8.3)	(11.1)	(2.8)	(5.6)	(8.3)	(11.1)		(5.6)	(8.3)	(8.3)						(2.8)	(100.0)				
腎・尿路結核	2	1	2	1	2		1		3	3		2	1								18		18	140.89	100.10
(%)	(11.1)	(5.6)	(11.1)	(5.6)	(11.1)		(5.6)		(16.7)	(16.7)		(11.1)	(5.6)								(100.0)				
腎・尿路結石	2	2	3	3	4	1	10	5	2	8		4	5	1	1		1	1	1	1	55	3	58	205.76	183.22
(%)	(3.8)	(3.6)	(5.5)	(5.5)	(7.3)	(1.8)	(18.2)	(9.1)	(3.6)	(14.5)		(7.3)	(9.1)	(1.8)	(1.8)		(1.8)	(1.8)	(1.8)	(1.8)	(100.0)				
腎・尿路腫瘍	7	3	4	2	10	7	9	5	5	9		6	2	1	1	1				1	73	2	75	153.60	133.98
(%)	(9.6)	(4.1)	(5.5)	(2.7)	(13.7)	(9.6)	(12.3)	(6.8)	(6.8)	(12.3)		(8.2)	(2.7)	(1.4)	(1.4)	(1.4)				(1.4)	(100.0)				
閉塞性尿路障害	2	2	4	4	4	4	3	3	1	7		7	6	2	1	2	1	1	1	3	58	3	61	273.90	327.71
(%)	(3.4)	(3.4)	(6.9)	(6.9)	(6.9)	(6.9)	(5.2)	(5.2)	(1.7)	(12.1)		(12.1)	(10.3)	(3.4)	(1.7)	(3.4)	(1.7)	(1.7)	(1.7)	(5.2)	(100.0)				
骨髄腫	1	1	3		1		2	2		2		3							1		16		16	174.63	162.98
(%)	(6.3)	(6.3)	(18.8)		(6.3)		(12.5)	(12.5)		(12.5)		(18.8)							(6.3)		(100.0)				
腎形成不全	3	5	5	7	10	6	12	3	6	8		12	4	4			2	2		1	90		90	189.10	211.56
(%)	(3.3)	(5.6)	(5.6)	(7.8)	(11.1)	(6.7)	(13.3)	(3.3)	(6.7)	(8.9)		(13.3)	(4.4)	(4.4)			(2.2)	(2.2)		(1.1)	(100.0)				
不明	129	193	227	247	257	278	210	169	184	412	5	236	141	77	50	9	19	22	30	32	2,927	66	2,993	171.28	157.28
(%)	(4.4)	(6.6)	(7.8)	(8.4)	(8.8)	(9.5)	(7.2)	(5.8)	(6.3)	(14.1)	(0.2)	(8.1)	(4.8)	(2.6)	(1.7)	(0.3)	(0.6)	(0.8)	(1.0)	(1.1)	(100.0)				
再導入	17	26	23	24	30	34	26	24	25	53		27	19	9	11		1	1	9	4	363	5	368	180.44	166.77
(%)	(4.7)	(7.2)	(6.3)	(6.6)	(8.3)	(9.4)	(7.2)	(6.6)	(6.9)	(14.6)		(7.4)	(5.2)	(2.5)	(3.0)		(0.3)	(0.3)	(2.5)	(1.1)	(100.0)				
その他	38	40	57	58	71	65	51	51	51	114	1	63	42	19	16	5	8	7	10	4	771	8	779	175.07	145.12
(%)	(4.9)	(5.2)	(7.4)	(7.5)	(9.2)	(8.4)	(6.6)	(6.6)	(6.6)	(14.8)	(0.1)	(8.2)	(5.4)	(2.5)	(2.1)	(0.6)	(1.0)	(0.9)	(1.3)	(0.5)	(100.0)				
合計	1,683	2,195	2,437	2,592	2,846	2,795	2,581	2,323	2,150	5,005	77	2,955	1,714	1,042	591	146	237	252	423	350	34,394	734	35,128	177.26	169.45
(%)	(4.9)	(6.4)	(7.1)	(7.5)	(8.3)	(8.1)	(7.5)	(6.8)	(6.3)	(14.6)	(0.2)	(8.6)	(5.0)	(3.0)	(1.7)	(0.4)	(0.7)	(0.7)	(1.2)	(1.0)	(100.0)				
記載なし																									
(%)																									
総計	1,683	2,195	2,437	2,592	2,846	2,795	2,581	2,323	2,150	5,005	77	2,955	1,714	1,042	591	146	237	252	423	350	34,394	734	35,128	177.26	169.45
(%)	(4.9)	(6.4)	(7.1)	(7.5)	(8.3)	(8.1)	(7.5)	(6.8)	(6.3)	(14.6)	(0.2)	(8.6)	(5.0)	(3.0)	(1.7)	(0.4)	(0.7)	(0.7)	(1.2)	(1.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy
2014年調査