

表 3534 PTH値(pg/mL) 透析後体重(kg)別（血液透析濾過患者 透析回数週3回）PTH測定法:intact-PTH

透析後体重	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
20未満					1				1			4	3					1			10				
(%)					(10.0)				(10.0)			(40.0)	(30.0)					(10.0)			(100.0)				
20～	8	7	5	6	3	4	5	1	4	4	1	5	5	3			2			2	65	2	67	175.32	200.16
(%)	(12.3)	(10.8)	(7.7)	(9.2)	(4.6)	(6.2)	(7.7)	(1.5)	(6.2)	(6.2)	(1.5)	(7.7)	(7.7)	(4.6)			(3.1)			(3.1)	(100.0)				
30～	156	197	193	177	216	182	177	148	124	311		178	91	52	44	9	16	11	28	33	2,343	55	2,398	168.99	195.37
(%)	(6.7)	(8.4)	(8.2)	(7.6)	(9.2)	(7.8)	(7.6)	(6.3)	(5.3)	(13.3)		(7.6)	(3.9)	(2.2)	(1.9)	(0.4)	(0.7)	(0.5)	(1.2)	(1.4)	(100.0)				
40～	541	620	694	643	771	745	631	579	503	1,196	21	662	400	266	145	30	57	68	85	82	8,739	162	8,901	169.23	173.67
(%)	(6.2)	(7.1)	(7.9)	(7.4)	(8.8)	(8.5)	(7.2)	(6.6)	(5.8)	(13.7)	(0.2)	(7.6)	(4.6)	(3.0)	(1.7)	(0.3)	(0.7)	(0.8)	(1.0)	(0.9)	(100.0)				
50～	550	725	785	888	846	887	847	739	729	1,545	27	906	523	324	188	57	62	74	143	97	10,942	199	11,141	173.86	161.94
(%)	(5.0)	(6.6)	(7.2)	(8.1)	(7.7)	(8.1)	(7.7)	(6.8)	(6.7)	(14.1)	(0.2)	(8.3)	(4.8)	(3.0)	(1.7)	(0.5)	(0.6)	(0.7)	(1.3)	(0.9)	(100.0)				
60～	261	405	492	540	621	565	577	519	452	1,157	19	708	400	213	113	23	50	58	94	68	7,335	112	7,447	181.61	163.09
(%)	(3.6)	(5.5)	(6.7)	(7.4)	(8.5)	(7.7)	(7.9)	(7.1)	(6.2)	(15.8)	(0.3)	(9.7)	(5.5)	(2.9)	(1.5)	(0.3)	(0.7)	(0.8)	(1.3)	(0.9)	(100.0)				
70～	107	164	168	219	229	269	209	210	202	477	8	301	171	104	54	18	29	23	37	34	3,033	38	3,071	188.72	160.24
(%)	(3.5)	(5.4)	(5.5)	(7.2)	(7.6)	(8.9)	(6.9)	(6.9)	(6.7)	(15.7)	(0.3)	(9.9)	(5.6)	(3.4)	(1.8)	(0.6)	(1.0)	(0.8)	(1.2)	(1.1)	(100.0)				
80～	32	42	50	83	98	76	72	67	88	188		102	78	56	25	5	14	10	22	13	1,121	11	1,132	207.39	187.41
(%)	(2.9)	(3.7)	(4.5)	(7.4)	(8.7)	(6.8)	(6.4)	(6.0)	(7.9)	(16.8)		(9.1)	(7.0)	(5.0)	(2.2)	(0.4)	(1.2)	(0.9)	(2.0)	(1.2)	(100.0)				
90～	16	16	27	18	26	35	33	32	19	65		53	17	11	14	2	3	4	10	10	411	5	416	213.94	199.80
(%)	(3.9)	(3.9)	(6.6)	(4.4)	(6.3)	(8.5)	(8.0)	(7.8)	(4.6)	(15.8)		(12.9)	(4.1)	(2.7)	(3.4)	(0.5)	(0.7)	(1.0)	(2.4)	(2.4)	(100.0)				
100～	3	7	6	6	11	10	9	10	8	16	1	14	10	3	4	2	1	1	2	3	127	4	131	207.67	165.63
(%)	(2.4)	(5.5)	(4.7)	(4.7)	(8.7)	(7.9)	(7.1)	(7.9)	(6.3)	(12.6)	(0.8)	(11.0)	(7.9)	(2.4)	(3.1)	(1.6)	(0.8)	(0.8)	(1.6)	(2.4)	(100.0)				
110～	1	6	2	7	2	1	2	5	3	14		2	5	4	1			1		1	55	1	56	204.40	162.03
(%)	(1.8)		(10.9)	(3.6)	(12.7)	(3.6)	(1.8)	(9.1)	(5.5)	(25.5)		(3.6)	(9.1)	(7.3)	(1.8)			(1.8)		(1.8)	(100.0)				
120～		1	2	1	2	3	5		1	6		4	2	1					2	2	32		32	257.28	244.28
(%)		(3.1)	(6.3)	(3.1)	(6.3)	(9.4)	(15.6)		(3.1)	(18.8)		(12.5)	(6.3)	(3.1)					(6.3)	(6.3)	(100.0)				
合計	1,675	2,184	2,428	2,583	2,831	2,778	2,566	2,310	2,134	4,979	77	2,939	1,705	1,037	588	146	234	251	423	345	34,213	590	34,803	177.19	169.36
(%)	(4.9)	(6.4)	(7.1)	(7.5)	(8.3)	(8.1)	(7.5)	(6.8)	(6.2)	(14.6)	(0.2)	(8.6)	(5.0)	(3.0)	(1.7)	(0.4)	(0.7)	(0.7)	(1.2)	(1.0)	(100.0)				
記載なし	8	11	9	9	15	17	15	13	16	26		16	9	5	3		3	1		5	181	144	325	190.03	185.65
(%)	(4.4)	(6.1)	(5.0)	(5.0)	(8.3)	(9.4)	(8.3)	(7.2)	(8.8)	(14.4)		(8.8)	(5.0)	(2.8)	(1.7)		(1.7)	(0.6)		(2.8)	(100.0)				
総計	1,683	2,195	2,437	2,592	2,846	2,795	2,581	2,323	2,150	5,005	77	2,955	1,714	1,042	591	146	237	252	423	350	34,394	734	35,128	177.26	169.45
(%)	(4.9)	(6.4)	(7.1)	(7.5)	(8.3)	(8.1)	(7.5)	(6.8)	(6.3)	(14.6)	(0.2)	(8.6)	(5.0)	(3.0)	(1.7)	(0.4)	(0.7)	(0.7)	(1.2)	(1.0)	(100.0)				
平均	53.69	54.58	55.24	56.16	56.27	56.60	56.60	56.88	57.30	57.63	57.45	57.99	58.09	57.82	57.32	57.78	58.21	57.49	58.61	57.60	56.69	54.46	56.65		
標準偏差	11.94	12.06	12.68	12.36	13.11	12.97	12.85	12.84	12.86	13.33	11.12	13.43	13.63	13.56	14.01	13.73	14.26	13.53	14.29	15.97	13.07	12.56	13.06		

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査