

表 3535 PTH値(pg/mL) 透析前ヘモグロビン濃度(g/dL)別（血液透析濾過患者 透析回数週3回）PTH測定法:intact-PTH

透析前ヘモグロビン濃度	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
5未満	2	3	3	2	3	1		2	2	3		1	1					1			24	1	25	135.46	126.62
(%)	(8.3)	(12.5)	(12.5)	(8.3)	(12.5)	(4.2)		(8.3)	(8.3)	(12.5)		(4.2)	(4.2)					(4.2)			(100.0)				
5～	2	2	3	4	3	2	2	1		3		2		1	1						26	2	28	132.23	118.07
(%)	(7.7)	(7.7)	(11.5)	(15.4)	(11.5)	(7.7)	(7.7)	(3.8)		(11.5)		(7.7)		(3.8)	(3.8)						(100.0)				
6～	5	5	8	6	12	5	6	11	7	12		5	5	3	1				2	1	94	6	100	162.64	140.75
(%)	(5.3)	(5.3)	(8.5)	(6.4)	(12.8)	(5.3)	(6.4)	(11.7)	(7.4)	(12.8)		(5.3)	(5.3)	(3.2)	(1.1)				(2.1)	(1.1)	(100.0)				
7～	24	35	36	28	38	36	24	29	18	46		23	12	10	8	1	2	1	8	10	389	20	409	194.94	335.12
(%)	(6.2)	(9.0)	(9.3)	(7.2)	(9.8)	(9.3)	(6.2)	(7.5)	(4.6)	(11.8)		(5.9)	(3.1)	(2.6)	(2.1)	(0.3)	(0.5)	(0.3)	(2.1)	(2.6)	(100.0)				
8～	103	126	114	126	100	118	86	97	72	175	4	109	69	46	25	7	11	15	15	15	1,433	32	1,465	171.45	195.04
(%)	(7.2)	(8.8)	(8.0)	(8.8)	(7.0)	(8.2)	(6.0)	(6.8)	(5.0)	(12.2)	(0.3)	(7.6)	(4.8)	(3.2)	(1.7)	(0.5)	(0.8)	(1.0)	(1.0)	(1.0)	(100.0)				
9～	265	364	410	410	399	405	378	343	325	686	11	418	265	142	84	28	43	36	78	64	5,154	111	5,265	177.27	172.11
(%)	(5.1)	(7.1)	(8.0)	(8.0)	(7.7)	(7.9)	(7.3)	(6.7)	(6.3)	(13.3)	(0.2)	(8.1)	(5.1)	(2.8)	(1.6)	(0.5)	(0.8)	(0.7)	(1.5)	(1.2)	(100.0)				
10～	489	694	810	868	1,003	899	841	804	706	1,767	29	1,026	573	360	194	49	70	80	158	111	11,531	176	11,707	178.60	161.62
(%)	(4.2)	(6.0)	(7.0)	(7.5)	(8.7)	(7.8)	(7.3)	(7.0)	(6.1)	(15.3)	(0.3)	(8.9)	(5.0)	(3.1)	(1.7)	(0.4)	(0.6)	(0.7)	(1.4)	(1.0)	(100.0)				
11～	495	607	688	752	857	867	827	664	696	1,506	23	878	508	295	172	37	67	76	106	82	10,203	126	10,329	174.35	159.40
(%)	(4.9)	(5.9)	(6.7)	(7.4)	(8.4)	(8.5)	(8.1)	(6.5)	(6.8)	(14.8)	(0.2)	(8.6)	(5.0)	(2.9)	(1.7)	(0.4)	(0.7)	(0.7)	(1.0)	(0.8)	(100.0)				
12～	203	255	244	279	295	337	299	266	236	541	8	336	200	128	71	15	32	29	34	40	3,848	62	3,910	178.19	175.83
(%)	(5.3)	(6.6)	(6.3)	(7.3)	(7.7)	(8.8)	(7.8)	(6.9)	(6.1)	(14.1)	(0.2)	(8.7)	(5.2)	(3.3)	(1.8)	(0.4)	(0.8)	(0.8)	(0.9)	(1.0)	(100.0)				
13～	51	60	64	74	84	70	91	66	56	173	1	104	51	34	19	8	9	6	13	15	1,049	13	1,062	187.37	171.84
(%)	(4.9)	(5.7)	(6.1)	(7.1)	(8.0)	(6.7)	(8.7)	(6.3)	(5.3)	(16.5)	(0.1)	(9.9)	(4.9)	(3.2)	(1.8)	(0.8)	(0.9)	(0.6)	(1.2)	(1.4)	(100.0)				
14～	24	33	30	33	36	42	22	31	24	69		43	22	16	11	1	2	5	9	11	464	2	466	196.82	204.40
(%)	(5.2)	(7.1)	(6.5)	(7.1)	(7.8)	(9.1)	(4.7)	(6.7)	(5.2)	(14.9)		(9.3)	(4.7)	(3.4)	(2.4)	(0.2)	(0.4)	(1.1)	(1.9)	(2.4)	(100.0)				
合計	1,663	2,184	2,410	2,582	2,830	2,782	2,576	2,314	2,142	4,981	76	2,945	1,706	1,035	586	146	236	249	423	349	34,215	551	34,766	177.38	169.55
(%)	(4.9)	(6.4)	(7.0)	(7.5)	(8.3)	(8.1)	(7.5)	(6.8)	(6.3)	(14.6)	(0.2)	(8.6)	(5.0)	(3.0)	(1.7)	(0.4)	(0.7)	(0.7)	(1.2)	(1.0)	(100.0)				
記載なし	20	11	27	10	16	13	5	9	8	24	1	10	8	7	5		1	3		1	179	183	362	154.00	148.29
(%)	(11.2)	(6.1)	(15.1)	(5.6)	(8.9)	(7.3)	(2.8)	(5.0)	(4.5)	(13.4)	(0.6)	(5.6)	(4.5)	(3.9)	(2.8)		(0.6)	(1.7)		(0.6)	(100.0)				
総計	1,683	2,195	2,437	2,592	2,846	2,795	2,581	2,323	2,150	5,005	77	2,955	1,714	1,042	591	146	237	252	423	350	34,394	734	35,128	177.26	169.45
(%)	(4.9)	(6.4)	(7.1)	(7.5)	(8.3)	(8.1)	(7.5)	(6.8)	(6.3)	(14.6)	(0.2)	(8.6)	(5.0)	(3.0)	(1.7)	(0.4)	(0.7)	(0.7)	(1.2)	(1.0)	(100.0)				
平均	10.79	10.76	10.75	10.79	10.82	10.86	10.89	10.82	10.87	10.89	10.74	10.90	10.87	10.88	10.89	10.84	10.86	10.85	10.70	10.81	10.84	10.51	10.83		
標準偏差	1.36	1.33	1.29	1.28	1.26	1.24	1.21	1.28	1.20	1.22	1.05	1.24	1.22	1.28	1.30	1.29	1.24	1.29	1.30	1.49	1.26	1.41	1.26		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査