

表 3536 PTH値(pg/mL) 総コレステロール濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）PTH測定法:intact-PTH

総コレステロール濃度	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
100未満	62	64	72	89	74	80	65	66	56	115	1	65	30	21	8	1	5	1	4	7	886	19	905	151.28	163.97
(%)	(7.0)	(7.2)	(8.1)	(10.0)	(8.4)	(9.0)	(7.3)	(7.4)	(6.3)	(13.0)	(0.1)	(7.3)	(3.4)	(2.4)	(0.9)	(0.1)	(0.6)	(0.1)	(0.5)	(0.8)	(100.0)				
100～	130	193	192	229	235	215	204	195	182	373	5	208	119	68	46	9	14	14	28	20	2,679	45	2,724	166.36	159.75
(%)	(4.9)	(7.2)	(7.2)	(8.5)	(8.8)	(8.0)	(7.6)	(7.3)	(6.8)	(13.9)	(0.2)	(7.8)	(4.4)	(2.5)	(1.7)	(0.3)	(0.5)	(0.5)	(1.0)	(0.7)	(100.0)				
120～	229	364	371	379	467	423	383	371	341	754	5	446	216	122	69	20	26	38	51	38	5,113	53	5,166	167.36	147.60
(%)	(4.5)	(7.1)	(7.3)	(7.4)	(9.1)	(8.3)	(7.5)	(7.3)	(6.7)	(14.7)	(0.1)	(8.7)	(4.2)	(2.4)	(1.3)	(0.4)	(0.5)	(0.7)	(1.0)	(0.7)	(100.0)				
140～	286	403	449	436	503	508	492	407	398	917	14	548	307	186	101	26	52	38	67	47	6,185	89	6,274	175.22	166.37
(%)	(4.6)	(6.5)	(7.3)	(7.0)	(8.1)	(8.2)	(8.0)	(6.6)	(6.4)	(14.8)	(0.2)	(8.9)	(5.0)	(3.0)	(1.6)	(0.4)	(0.8)	(0.6)	(1.1)	(0.8)	(100.0)				
160～	262	312	393	409	426	430	409	315	330	791	19	477	278	164	103	17	44	42	59	53	5,333	80	5,413	178.71	167.25
(%)	(4.9)	(5.9)	(7.4)	(7.7)	(8.0)	(8.1)	(7.7)	(5.9)	(6.2)	(14.8)	(0.4)	(8.9)	(5.2)	(3.1)	(1.9)	(0.3)	(0.8)	(0.8)	(1.1)	(1.0)	(100.0)				
180～	163	209	224	254	253	300	250	230	201	511	8	302	170	127	68	16	29	29	48	52	3,444	52	3,496	188.08	183.05
(%)	(4.7)	(6.1)	(6.5)	(7.4)	(7.3)	(8.7)	(7.3)	(6.7)	(5.8)	(14.8)	(0.2)	(8.8)	(4.9)	(3.7)	(2.0)	(0.5)	(0.8)	(0.8)	(1.4)	(1.5)	(100.0)				
200～	94	112	135	143	144	165	140	130	120	268	2	202	113	72	47	17	14	14	31	26	1,989	24	2,013	190.40	174.45
(%)	(4.7)	(5.6)	(6.8)	(7.2)	(7.2)	(8.3)	(7.0)	(6.5)	(6.0)	(13.5)	(0.1)	(10.2)	(5.7)	(3.6)	(2.4)	(0.9)	(0.7)	(0.7)	(1.6)	(1.3)	(100.0)				
220～	53	46	50	54	81	70	68	52	36	124	1	77	55	23	18	7	3	3	22	9	852	11	863	186.19	169.43
(%)	(6.2)	(5.4)	(5.9)	(6.3)	(9.5)	(8.2)	(8.0)	(6.1)	(4.2)	(14.6)	(0.1)	(9.0)	(6.5)	(2.7)	(2.1)	(0.8)	(0.4)	(0.4)	(2.6)	(1.1)	(100.0)				
240～	17	27	25	29	28	23	20	20	18	32	2	47	17	19	6	1	1	2	5	3	342	2	344	181.52	159.07
(%)	(5.0)	(7.9)	(7.3)	(8.5)	(8.2)	(6.7)	(5.8)	(5.8)	(5.3)	(9.4)	(0.6)	(13.7)	(5.0)	(5.6)	(1.8)	(0.3)	(0.3)	(0.6)	(1.5)	(0.9)	(100.0)				
260～	12	7	10	19	9	14	10	14	15	27		13	13	11	11	2	3	2	3	4	199	2	201	229.89	305.54
(%)	(6.0)	(3.5)	(5.0)	(9.5)	(4.5)	(7.0)	(5.0)	(7.0)	(7.5)	(13.6)		(6.5)	(6.5)	(5.5)	(5.5)	(1.0)	(1.5)	(1.0)	(1.5)	(2.0)	(100.0)				
合計	1,308	1,737	1,921	2,041	2,220	2,228	2,041	1,800	1,697	3,912	57	2,385	1,318	813	477	116	191	183	318	259	27,022	377	27,399	176.34	167.05
(%)	(4.8)	(6.4)	(7.1)	(7.6)	(8.2)	(8.2)	(7.6)	(6.7)	(6.3)	(14.5)	(0.2)	(8.8)	(4.9)	(3.0)	(1.8)	(0.4)	(0.7)	(0.7)	(1.2)	(1.0)	(100.0)				
記載なし	375	458	516	551	626	567	540	523	453	1,093	20	570	396	229	114	30	46	69	105	91	7,372	357	7,729	180.61	177.97
(%)	(5.1)	(6.2)	(7.0)	(7.5)	(8.5)	(7.7)	(7.3)	(7.1)	(6.1)	(14.8)	(0.3)	(7.7)	(5.4)	(3.1)	(1.5)	(0.4)	(0.6)	(0.9)	(1.4)	(1.2)	(100.0)				
総計	1,683	2,195	2,437	2,592	2,846	2,795	2,581	2,323	2,150	5,005	77	2,955	1,714	1,042	591	146	237	252	423	350	34,394	734	35,128	177.26	169.45
(%)	(4.9)	(6.4)	(7.1)	(7.5)	(8.3)	(8.1)	(7.5)	(6.8)	(6.3)	(14.6)	(0.2)	(8.6)	(5.0)	(3.0)	(1.7)	(0.4)	(0.7)	(0.7)	(1.2)	(1.0)	(100.0)				
平均	157.78	155.27	156.16	156.43	155.73	157.48	157.34	156.35	156.16	157.47	163.00	160.20	161.44	163.32	164.94	166.99	161.42	161.09	165.94	165.41	157.89	155.92	157.86		
標準偏差	37.69	35.47	35.56	36.92	35.61	35.22	34.98	36.08	35.32	34.53	32.05	35.68	35.82	37.14	37.15	38.36	34.10	32.13	37.09	36.68	35.75	35.76	35.75		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy
2014年調査