

表 3537 PTH値(pg/mL) HDL-C濃度(mg/dL)別 (血液透析濾過患者 透析回数週3回) PTH測定法:intact-PTH

HDL-C濃度	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
20未満	11	9	8	3	7	3	6		3	11		4	3	2	2	2			2	1	77		77	158.39	167.01
(%)	(14.3)	(11.7)	(10.4)	(3.9)	(9.1)	(3.9)	(7.8)		(3.9)	(14.3)		(5.2)	(3.9)	(2.6)	(2.6)	(2.6)			(2.6)	(1.3)	(100.0)				
20～	91	114	117	152	150	139	134	105	91	251	2	149	82	44	22	3	7	11	8	14	1,686	24	1,710	163.14	144.95
(%)	(5.4)	(6.8)	(6.9)	(9.0)	(8.9)	(8.2)	(7.9)	(6.2)	(5.4)	(14.9)	(0.1)	(8.8)	(4.9)	(2.6)	(1.3)	(0.2)	(0.4)	(0.7)	(0.5)	(0.8)	(100.0)				
30～	280	338	386	425	445	461	406	379	356	791	10	473	260	169	96	15	37	43	66	54	5,490	68	5,558	175.77	164.22
(%)	(5.1)	(6.2)	(7.0)	(7.7)	(8.1)	(8.4)	(7.4)	(6.9)	(6.5)	(14.4)	(0.2)	(8.6)	(4.7)	(3.1)	(1.7)	(0.3)	(0.7)	(0.8)	(1.2)	(1.0)	(100.0)				
40～	332	417	464	476	543	538	504	443	446	967	11	528	279	204	107	50	39	56	81	57	6,542	107	6,649	174.76	159.41
(%)	(5.1)	(6.4)	(7.1)	(7.3)	(8.3)	(8.2)	(7.7)	(6.8)	(6.8)	(14.8)	(0.2)	(8.1)	(4.3)	(3.1)	(1.6)	(0.8)	(0.6)	(0.9)	(1.2)	(0.9)	(100.0)				
50～	235	349	381	413	432	417	403	343	327	761	14	459	269	176	92	11	49	29	73	56	5,289	73	5,362	180.00	180.88
(%)	(4.4)	(6.6)	(7.2)	(7.8)	(8.2)	(7.9)	(7.6)	(6.5)	(6.2)	(14.4)	(0.3)	(8.7)	(5.1)	(3.3)	(1.7)	(0.2)	(0.9)	(0.5)	(1.4)	(1.1)	(100.0)				
60～	147	199	272	244	273	278	261	241	223	511	10	292	184	95	57	13	21	27	30	40	3,418	42	3,460	177.92	167.37
(%)	(4.3)	(5.8)	(8.0)	(7.1)	(8.0)	(8.1)	(7.6)	(7.1)	(6.5)	(15.0)	(0.3)	(8.5)	(5.4)	(2.8)	(1.7)	(0.4)	(0.6)	(0.8)	(0.9)	(1.2)	(100.0)				
70～	88	110	103	131	148	154	143	116	113	254	5	156	103	58	36	8	12	15	16	23	1,792	18	1,810	184.48	191.35
(%)	(4.9)	(6.1)	(5.7)	(7.3)	(8.3)	(8.6)	(8.0)	(6.5)	(6.3)	(14.2)	(0.3)	(8.7)	(5.7)	(3.2)	(2.0)	(0.4)	(0.7)	(0.8)	(0.9)	(1.3)	(100.0)				
80～	42	42	69	72	74	58	81	67	52	110	1	95	39	28	22	6	5	5	15	9	892	8	900	185.85	205.17
(%)	(4.7)	(4.7)	(7.7)	(8.1)	(8.3)	(6.5)	(9.1)	(7.5)	(5.8)	(12.3)	(0.1)	(10.7)	(4.4)	(3.1)	(2.5)	(0.7)	(0.6)	(0.6)	(1.7)	(1.0)	(100.0)				
90～	15	26	28	19	34	28	32	20	25	65	2	38	17	15	8	1	3	5	5	1	387	5	392	179.68	138.66
(%)	(3.9)	(6.7)	(7.2)	(4.9)	(8.8)	(7.2)	(8.3)	(5.2)	(6.5)	(16.8)	(0.5)	(9.8)	(4.4)	(3.9)	(2.1)	(0.3)	(0.8)	(1.3)	(1.3)	(0.3)	(100.0)				
100～	18	30	21	22	25	31	24	16	22	47		33	16	8	5		3	1	5	2	329	3	332	168.34	144.72
(%)	(5.5)	(9.1)	(6.4)	(6.7)	(7.6)	(9.4)	(7.3)	(4.9)	(6.7)	(14.3)		(10.0)	(4.9)	(2.4)	(1.5)		(0.9)	(0.3)	(1.5)	(0.6)	(100.0)				
合計	1,259	1,634	1,849	1,957	2,131	2,107	1,994	1,730	1,658	3,768	55	2,227	1,252	799	447	109	176	192	301	257	25,902	348	26,250	176.70	168.86
(%)	(4.9)	(6.3)	(7.1)	(7.6)	(8.2)	(8.1)	(7.7)	(6.7)	(6.4)	(14.5)	(0.2)	(8.6)	(4.8)	(3.1)	(1.7)	(0.4)	(0.7)	(0.7)	(1.2)	(1.0)	(100.0)				
記載なし	424	561	588	635	715	688	587	593	492	1,237	22	728	462	243	144	37	61	60	122	93	8,492	386	8,878	178.95	171.26
(%)	(5.0)	(6.6)	(6.9)	(7.5)	(8.4)	(8.1)	(6.9)	(7.0)	(5.8)	(14.6)	(0.3)	(8.6)	(5.4)	(2.9)	(1.7)	(0.4)	(0.7)	(0.7)	(1.4)	(1.1)	(100.0)				
総計	1,683	2,195	2,437	2,592	2,846	2,795	2,581	2,323	2,150	5,005	77	2,955	1,714	1,042	591	146	237	252	423	350	34,394	734	35,128	177.26	169.45
(%)	(4.9)	(6.4)	(7.1)	(7.5)	(8.3)	(8.1)	(7.5)	(6.8)	(6.3)	(14.6)	(0.2)	(8.6)	(5.0)	(3.0)	(1.7)	(0.4)	(0.7)	(0.7)	(1.2)	(1.0)	(100.0)				
平均	49.73	50.58	50.84	50.12	50.68	50.52	51.04	50.71	50.86	50.66	53.02	51.37	51.42	51.10	52.00	50.14	51.93	50.37	51.43	51.56	50.78	49.26	50.76		
標準偏差	17.20	17.71	17.16	16.73	17.23	17.07	17.13	16.67	16.93	17.12	15.33	17.76	17.10	17.71	17.71	15.31	18.06	16.40	17.45	16.14	17.17	15.89	17.15		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査