

表 3538 PTH値(pg/mL) Non-HDL-C濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）PTH測定法:intact-PTH

Non-HDL-C濃度	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
50未満	26	29	42	39	41	50	46	41	25	64	1	42	15	10	5			1	1	4	482	6	488	150.61	124.96
(%)	(5.4)	(6.0)	(8.7)	(8.1)	(8.5)	(10.4)	(9.5)	(8.5)	(5.2)	(13.3)	(0.2)	(8.7)	(3.1)	(2.1)	(1.0)			(0.2)	(0.2)	(0.8)	(100.0)				
50～	100	162	150	178	182	190	182	150	148	310	2	172	87	53	30	7	11	11	24	18	2,167	18	2,185	167.40	175.13
(%)	(4.6)	(7.5)	(6.9)	(8.2)	(8.4)	(8.8)	(8.4)	(6.9)	(6.8)	(14.3)	(0.1)	(7.9)	(4.0)	(2.4)	(1.4)	(0.3)	(0.5)	(0.5)	(1.1)	(0.8)	(100.0)				
70～	196	309	337	356	407	364	340	329	322	670	10	395	215	131	71	21	28	31	49	33	4,614	40	4,654	171.27	154.99
(%)	(4.2)	(6.7)	(7.3)	(7.7)	(8.8)	(7.9)	(7.4)	(7.1)	(7.0)	(14.5)	(0.2)	(8.6)	(4.7)	(2.8)	(1.5)	(0.5)	(0.6)	(0.7)	(1.1)	(0.7)	(100.0)				
90～	282	354	415	393	476	418	434	371	356	835	10	488	306	169	88	20	48	44	55	37	5,599	67	5,666	173.59	150.70
(%)	(5.0)	(6.3)	(7.4)	(7.0)	(8.5)	(7.5)	(7.8)	(6.6)	(6.4)	(14.9)	(0.2)	(8.7)	(5.5)	(3.0)	(1.6)	(0.4)	(0.9)	(0.8)	(1.0)	(0.7)	(100.0)				
110～	196	287	312	357	361	388	367	281	289	697	16	404	201	141	76	18	34	37	55	54	4,571	71	4,642	180.05	175.74
(%)	(4.3)	(6.3)	(6.8)	(7.8)	(7.9)	(8.5)	(8.0)	(6.1)	(6.3)	(15.2)	(0.4)	(8.8)	(4.4)	(3.1)	(1.7)	(0.4)	(0.7)	(0.8)	(1.2)	(1.2)	(100.0)				
130～	165	180	191	195	213	257	206	180	160	380	5	260	121	94	66	12	22	19	37	34	2,797	37	2,834	180.19	173.70
(%)	(5.9)	(6.4)	(6.8)	(7.0)	(7.6)	(9.2)	(7.4)	(6.4)	(5.7)	(13.6)	(0.2)	(9.3)	(4.3)	(3.4)	(2.4)	(0.4)	(0.8)	(0.7)	(1.3)	(1.2)	(100.0)				
150～	68	70	94	89	117	112	106	90	78	197	4	142	97	52	36	10	13	11	17	28	1,431	13	1,444	199.52	193.49
(%)	(4.8)	(4.9)	(6.6)	(6.2)	(8.2)	(7.8)	(7.4)	(6.3)	(5.5)	(13.8)	(0.3)	(9.9)	(6.8)	(3.6)	(2.5)	(0.7)	(0.9)	(0.8)	(1.2)	(2.0)	(100.0)				
170～	35	35	40	52	45	52	35	42	41	86		54	36	21	17	6	1	4	7	6	615	5	620	185.07	205.55
(%)	(5.7)	(5.7)	(6.5)	(8.5)	(7.3)	(8.5)	(5.7)	(6.8)	(6.7)	(14.0)		(8.8)	(5.9)	(3.4)	(2.8)	(1.0)	(0.2)	(0.7)	(1.1)	(1.0)	(100.0)				
190～	10	11	11	20	18	12	17	16	13	39		21	14	10	7	2	1	3	9	1	235	4	239	202.96	168.55
(%)	(4.3)	(4.7)	(4.7)	(8.5)	(7.7)	(5.1)	(7.2)	(6.8)	(5.5)	(16.6)		(8.9)	(6.0)	(4.3)	(3.0)	(0.9)	(0.4)	(1.3)	(3.8)	(0.4)	(100.0)				
210～	8	7	11	15	6	11	10	7	9	10	1	8	6	9	3	1	3	1	3	2	131	1	132	193.86	177.66
(%)	(6.1)	(5.3)	(8.4)	(11.5)	(4.6)	(8.4)	(7.6)	(5.3)	(6.9)	(7.6)	(0.8)	(6.1)	(4.6)	(6.9)	(2.3)	(0.8)	(2.3)	(0.8)	(2.3)	(1.5)	(100.0)				
合計	1,086	1,444	1,603	1,694	1,866	1,854	1,743	1,507	1,441	3,288	49	1,986	1,098	690	399	97	161	162	257	217	22,642	262	22,904	176.53	166.73
(%)	(4.8)	(6.4)	(7.1)	(7.5)	(8.2)	(8.2)	(7.7)	(6.7)	(6.4)	(14.5)	(0.2)	(8.8)	(4.8)	(3.0)	(1.8)	(0.4)	(0.7)	(0.7)	(1.1)	(1.0)	(100.0)				
記載なし	597	751	834	898	980	941	838	816	709	1,717	28	969	616	352	192	49	76	90	166	133	11,752	472	12,224	178.66	174.58
(%)	(5.1)	(6.4)	(7.1)	(7.6)	(8.3)	(8.0)	(7.1)	(6.9)	(6.0)	(14.6)	(0.2)	(8.2)	(5.2)	(3.0)	(1.6)	(0.4)	(0.6)	(0.8)	(1.4)	(1.1)	(100.0)				
総計	1,683	2,195	2,437	2,592	2,846	2,795	2,581	2,323	2,150	5,005	77	2,955	1,714	1,042	591	146	237	252	423	350	34,394	734	35,128	177.26	169.45
(%)	(4.9)	(6.4)	(7.1)	(7.5)	(8.3)	(8.1)	(7.5)	(6.8)	(6.3)	(14.6)	(0.2)	(8.6)	(5.0)	(3.0)	(1.7)	(0.4)	(0.7)	(0.7)	(1.2)	(1.0)	(100.0)				
平均	108.58	104.85	105.87	106.32	105.27	106.80	106.01	105.54	105.47	106.76	111.02	107.95	109.57	110.91	113.96	114.58	111.25	111.23	112.94	114.34	107.03	109.72	107.06		
標準偏差	34.03	32.62	33.23	34.61	32.80	33.23	33.89	34.22	33.00	32.45	32.71	33.52	33.64	35.19	34.14	36.85	32.22	32.12	36.25	35.01	33.52	31.72	33.50		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査