

表 3545 PTH値(pg/mL) 原疾患別（腹膜透析患者）PTH測定法intact-PTH

原疾患	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	33	98	95	111	113	103	104	103	102	229	5	157	107	74	47	16	19	30	47	48	1,641	135	1,776	237.17	258.57
(%)	(2.0)	(6.0)	(5.8)	(6.8)	(6.9)	(6.3)	(6.3)	(6.3)	(6.2)	(14.0)	(0.3)	(9.6)	(6.5)	(4.5)	(2.9)	(1.0)	(1.2)	(1.8)	(2.9)	(2.9)	(100.0)				
慢性腎盂腎炎	2	5	2	5	3	2	2	4	2	5		11	3	2	2			1	2	1	54	2	56	225.41	215.86
(%)	(3.7)	(9.3)	(3.7)	(9.3)	(5.6)	(3.7)	(3.7)	(7.4)	(3.7)	(9.3)		(20.4)	(5.6)	(3.7)	(3.7)			(1.9)	(3.7)	(1.9)	(100.0)				
急速進行性糸球体腎炎	1	3	1	1	2	2	4	5	1	3		4	2	4		1		1	2		37	4	41	230.32	177.55
(%)	(2.7)	(8.1)	(2.7)	(2.7)	(5.4)	(5.4)	(10.8)	(13.5)	(2.7)	(8.1)		(10.8)	(5.4)	(10.8)		(2.7)		(2.7)	(5.4)		(100.0)				
妊娠腎/妊娠中毒症	1			3	1	2			1	3		1	2						1	1	16	1	17	249.69	261.69
(%)	(6.3)			(18.8)	(6.3)	(12.5)			(6.3)	(18.8)		(6.3)	(12.5)						(6.3)	(6.3)	(100.0)				
その他の分類不能の腎炎		1	5	3	4	4	3	5	2	5		4	4	1	3	1			1	1	47	2	49	220.09	197.47
(%)		(2.1)	(10.6)	(6.4)	(8.5)	(8.5)	(6.4)	(10.6)	(4.3)	(10.6)		(8.5)	(8.5)	(2.1)	(6.4)	(2.1)			(2.1)	(2.1)	(100.0)				
多発性嚢胞腎	6	7	6	12	5	7	9	10	5	22		12	7	7	3	1	2	1	5	1	128	7	135	209.41	172.65
(%)	(4.7)	(5.5)	(4.7)	(9.4)	(3.9)	(5.5)	(7.0)	(7.8)	(3.9)	(17.2)		(9.4)	(5.5)	(5.5)	(2.3)	(0.8)	(1.6)	(0.8)	(3.9)	(0.8)	(100.0)				
腎硬化症	21	33	32	35	39	38	39	33	38	88	1	49	47	22	21	5	6	9	7	9	572	43	615	208.88	180.02
(%)	(3.7)	(5.8)	(5.6)	(6.1)	(6.8)	(6.6)	(6.8)	(5.8)	(6.6)	(15.4)	(0.2)	(8.6)	(8.2)	(3.8)	(3.7)	(0.9)	(1.0)	(1.6)	(1.2)	(1.6)	(100.0)				
悪性高血圧	1	3	5	2	2	6	7	4	4	8		4	2	1	2	1		1	1		54	1	55	183.52	134.17
(%)	(1.9)	(5.6)	(9.3)	(3.7)	(3.7)	(11.1)	(13.0)	(7.4)	(7.4)	(14.8)		(7.4)	(3.7)	(1.9)	(3.7)	(1.9)		(1.9)	(1.9)		(100.0)				
糖尿病性腎症	43	85	81	89	96	98	87	83	78	192	3	130	101	60	41	11	19	19	21	18	1,355	123	1,478	205.71	174.52
(%)	(3.2)	(6.3)	(6.0)	(6.6)	(7.1)	(7.2)	(6.4)	(6.1)	(5.8)	(14.2)	(0.2)	(9.6)	(7.5)	(4.4)	(3.0)	(0.8)	(1.4)	(1.4)	(1.5)	(1.3)	(100.0)				
SLE腎炎		1	3		1	2	1		2	4		3	2	2	2					2	25	6	31	314.12	397.28
(%)		(4.0)	(12.0)		(4.0)	(8.0)	(4.0)		(8.0)	(16.0)		(12.0)	(8.0)	(8.0)	(8.0)					(8.0)	(100.0)				
アミロイド腎						1				1											2	1	3	159.50	65.76
(%)						(50.0)				(50.0)											(100.0)				
痛風腎	1		1	1	1		1	2	3	4		2							1	1	18	1	19	252.28	290.60
(%)	(5.6)		(5.6)	(5.6)	(5.6)		(5.6)	(11.1)	(16.7)	(22.2)		(11.1)							(5.6)	(5.6)	(100.0)				
先天性代謝異常による腎不全					2				1			1		1			2		1		8	1	9	343.50	219.63
(%)					(25.0)				(12.5)			(12.5)		(12.5)			(25.0)		(12.5)		(100.0)				
腎・尿路結核				1																	1		1	64.00	
(%)				(100.0)																	(100.0)				
腎・尿路結石	1		1	1		1	1			4		1									10		10	162.60	101.86
(%)	(10.0)		(10.0)	(10.0)		(10.0)	(10.0)			(40.0)		(10.0)									(100.0)				
腎・尿路腫瘍	1		1		2									1						1	6	1	7	241.83	319.52
(%)	(16.7)		(16.7)		(33.3)									(16.7)						(16.7)	(100.0)				
閉塞性尿路障害		1				1	1		2	2		1		1			1	1	1	2	13		13	418.38	327.13
(%)		(7.7)				(25.0)	(25.0)		(15.4)	(15.4)		(7.7)		(7.7)			(7.7)	(7.7)	(7.7)	(15.4)	(100.0)				
骨髄腫					1		1	1											1		4		4	248.75	255.11
(%)					(25.0)		(25.0)	(25.0)											(25.0)		(100.0)				
腎形成不全	1	3	2	5	4	2	2	2	1	10		4	4	1	2		1	1		5	50		50	290.22	321.52
(%)	(2.0)	(6.0)	(4.0)	(10.0)	(8.0)	(4.0)	(4.0)	(4.0)	(2.0)	(20.0)		(8.0)	(8.0)	(2.0)	(4.0)		(2.0)	(2.0)		(10.0)	(100.0)				
不明	10	30	24	31	26	37	28	21	28	53	1	39	37	12	11	6	3	6	13	13	429	37	466	233.50	292.90
(%)	(2.3)	(7.0)	(5.6)	(7.2)	(6.1)	(8.6)	(6.5)	(4.9)	(6.5)	(12.4)	(0.2)	(9.1)	(8.6)	(2.8)	(2.6)	(1.4)	(0.7)	(1.4)	(3.0)	(3.0)	(100.0)				
再導入	1	2	4	3		1	2	3		7		3	1	1	1		1		1	2	32	3	35	249.50	273.68
(%)	(3.1)	(6.3)	(12.5)	(9.4)		(3.1)	(6.3)	(9.4)		(21.9)		(9.4)	(3.1)	(3.1)	(3.1)		(3.1)		(3.1)	(6.3)	(100.0)				
その他		9	5	12	7	10	13	9	6	14		10	8	10	4	1	2	1		5	126	8	134	231.98	265.44
(%)		(7.1)	(4.0)	(9.5)	(5.6)	(7.9)	(10.3)	(7.1)	(4.8)	(11.1)		(7.9)	(6.3)	(7.9)	(3.2)	(0.8)	(1.6)	(0.8)		(4.0)	(100.0)				
合計	123	281	268	315	309	316	305	285	276	654	10	431	331	200	139	43	54	73	105	110	4,628	376	5,004	223.88	229.31
(%)	(2.7)	(6.1)	(5.8)	(6.8)	(6.7)	(6.8)	(6.6)	(6.2)	(6.0)	(14.1)	(0.2)	(9.3)	(7.2)	(4.3)	(3.0)	(0.9)	(1.2)	(1.6)	(2.3)	(2.4)	(100.0)				
記載なし																									
(%)																									
総計	123	281	268	315	309	316	305	285	276	654	10	431	331	200	139	43	54	73	105	110	4,628	376	5,004	223.88	229.31
(%)	(2.7)	(6.1)	(5.8)	(6.8)	(6.7)	(6.8)	(6.6)	(6.2)	(6.0)	(14.1)	(0.2)	(9.3)	(7.2)	(4.3)	(3.0)	(0.9)	(1.2)	(1.6)	(2.3)	(2.4)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy
2014年調査