

表 3547 PTH値(pg/mL) 体重(kg)別（腹膜透析患者）PTH測定法:intact-PTH

体重	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
20未満	2	4	1	3	3	2		1	2	3		3		3	1				1	5	34		34	337.09	445.70
(%)	(5.9)	(11.8)	(2.9)	(8.8)	(8.8)	(5.9)		(2.9)	(5.9)	(8.8)		(8.8)		(8.8)	(2.9)				(2.9)	(14.7)	(100.0)				
20～	1	3	1		2	1	2		1	1		1	1	1	1				3	1	20	1	21	287.10	289.39
(%)	(5.0)	(15.0)	(5.0)		(10.0)	(5.0)	(10.0)		(5.0)	(5.0)		(5.0)	(5.0)	(5.0)	(5.0)				(15.0)	(5.0)	(100.0)				
30～	4	8	8	8	6	7	10	4	8	15		10	9	3	6	2	3		4	6	121	10	131	246.66	242.00
(%)	(3.3)	(6.6)	(6.6)	(6.6)	(5.0)	(5.8)	(8.3)	(3.3)	(6.6)	(12.4)		(8.3)	(7.4)	(2.5)	(5.0)	(1.7)	(2.5)		(3.3)	(5.0)	(100.0)				
40～	18	47	32	54	48	57	29	39	39	85	1	64	43	37	26	8	5	10	22	19	683	37	720	235.40	246.48
(%)	(2.6)	(6.9)	(4.7)	(7.9)	(7.0)	(8.3)	(4.2)	(5.7)	(5.7)	(12.4)	(0.1)	(9.4)	(6.3)	(5.4)	(3.8)	(1.2)	(0.7)	(1.5)	(3.2)	(2.8)	(100.0)				
50～	32	76	73	88	77	100	90	80	68	166	4	110	88	47	32	5	8	19	24	27	1,214	68	1,282	217.77	262.28
(%)	(2.6)	(6.3)	(6.0)	(7.2)	(6.3)	(8.2)	(7.4)	(6.6)	(5.6)	(13.7)	(0.3)	(9.1)	(7.2)	(3.9)	(2.6)	(0.4)	(0.7)	(1.6)	(2.0)	(2.2)	(100.0)				
60～	32	68	74	86	84	67	80	76	75	193	3	108	94	46	28	11	14	20	21	23	1,203	63	1,266	212.52	190.37
(%)	(2.7)	(5.7)	(6.2)	(7.1)	(7.0)	(5.6)	(6.7)	(6.3)	(6.2)	(16.0)	(0.2)	(9.0)	(7.8)	(3.8)	(2.3)	(0.9)	(1.2)	(1.7)	(1.7)	(1.9)	(100.0)				
70～	15	35	32	35	43	33	37	36	32	86		71	37	29	23	5	7	8	11	12	587	41	628	221.62	192.72
(%)	(2.6)	(6.0)	(5.5)	(6.0)	(7.3)	(5.6)	(6.3)	(6.1)	(5.5)	(14.7)		(12.1)	(6.3)	(4.9)	(3.9)	(0.9)	(1.2)	(1.4)	(1.9)	(2.0)	(100.0)				
80～	5	9	13	9	12	12	17	16	16	36		16	27	8	4	5	7	4	6	6	228	10	238	249.07	246.70
(%)	(2.2)	(3.9)	(5.7)	(3.9)	(5.3)	(5.3)	(7.5)	(7.0)	(7.0)	(15.8)		(7.0)	(11.8)	(3.5)	(1.8)	(2.2)	(3.1)	(1.8)	(2.6)	(2.6)	(100.0)				
90～	3	7	6	6	7	2	3	4	3	8		10	6	5	2	2	2	1	2		77	3	80	202.34	162.87
(%)	(3.9)	(9.1)	(7.8)	(7.8)	(9.1)	(2.6)	(3.9)	(5.2)	(3.9)	(10.4)		(13.0)	(7.8)	(6.5)		(2.6)	(2.6)	(1.3)	(2.6)		(100.0)				
100～		1	1			3	3	1	3					2	1		1				16	1	17	200.56	145.59
(%)		(6.3)	(6.3)			(18.8)	(18.8)	(6.3)	(18.8)					(12.5)	(6.3)		(6.3)				(100.0)				
110～	1		1			1			1		1	1						1		1	8		8	367.13	485.99
(%)	(12.5)		(12.5)			(12.5)			(12.5)		(12.5)	(12.5)						(12.5)		(12.5)	(100.0)				
120～							1		1				1		1						7	1	8	245.14	117.72
(%)						(14.3)	(14.3)		(14.3)	(42.9)			(14.3)		(14.3)						(100.0)				
合計	113	258	242	289	282	285	272	257	249	596	9	394	306	181	123	38	47	63	94	100	4,198	235	4,433	223.48	231.28
(%)	(2.7)	(6.1)	(5.8)	(6.9)	(6.7)	(6.8)	(6.5)	(6.1)	(5.9)	(14.2)	(0.2)	(9.4)	(7.3)	(4.3)	(2.9)	(0.9)	(1.1)	(1.5)	(2.2)	(2.4)	(100.0)				
記載なし	10	23	26	26	27	31	33	28	27	58	1	37	25	19	16	5	7	10	11	10	430	141	571	227.72	209.33
(%)	(2.3)	(5.3)	(6.0)	(6.0)	(6.3)	(7.2)	(7.7)	(6.5)	(6.3)	(13.5)	(0.2)	(8.6)	(5.8)	(4.4)	(3.7)	(1.2)	(1.6)	(2.3)	(2.6)	(2.3)	(100.0)				
総計	123	281	268	315	309	316	305	285	276	654	10	431	331	200	139	43	54	73	105	110	4,628	376	5,004	223.88	229.31
(%)	(2.7)	(6.1)	(5.8)	(6.8)	(6.7)	(6.8)	(6.6)	(6.2)	(6.0)	(14.1)	(0.2)	(9.3)	(7.2)	(4.3)	(3.0)	(0.9)	(1.2)	(1.6)	(2.3)	(2.4)	(100.0)				
平均	59.68	59.01	61.07	59.19	60.43	58.82	61.79	61.11	61.42	61.59	65.63	61.19	61.96	60.46	58.70	63.75	65.33	62.29	57.51	55.81	60.60	60.93	60.62		
標準偏差	15.39	14.49	13.65	12.84	13.40	13.43	13.78	12.45	15.84	13.60	18.59	13.34	14.03	15.21	14.71	15.99	16.60	13.24	14.90	17.28	14.08	13.51	14.05		

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査