

表 3550 PTH値(pg/mL) HDL-C濃度(mg/dL)別（腹膜透析患者）PTH測定法:intact-PTH

HDL-C濃度	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
20未満				1	1	1	1			1		1									6	1	7	142.67	71.81
(%)				(16.7)	(16.7)	(16.7)	(16.7)			(16.7)		(16.7)									(100.0)				
20～	4	20	19	13	12	24	17	9	10	35	1	17	21	12	4		2	3	3	1	227	8	235	187.34	146.94
(%)	(1.8)	(8.8)	(8.4)	(5.7)	(5.3)	(10.6)	(7.5)	(4.0)	(4.4)	(15.4)	(0.4)	(7.5)	(9.3)	(5.3)	(1.8)		(0.9)	(1.3)	(1.3)	(0.4)	(100.0)				
30～	30	46	43	53	52	46	52	58	53	94		81	43	31	18	5	9	12	14	12	752	22	774	204.55	187.40
(%)	(4.0)	(6.1)	(5.7)	(7.0)	(6.9)	(6.1)	(6.9)	(7.7)	(7.0)	(12.5)		(10.8)	(5.7)	(4.1)	(2.4)	(0.7)	(1.2)	(1.6)	(1.9)	(1.6)	(100.0)				
40～	24	46	47	46	56	55	49	54	52	110	2	75	54	30	26	12	8	14	9	20	789	13	802	227.43	272.93
(%)	(3.0)	(5.8)	(6.0)	(5.8)	(7.1)	(7.0)	(6.2)	(6.8)	(6.6)	(13.9)	(0.3)	(9.5)	(6.8)	(3.8)	(3.3)	(1.5)	(1.0)	(1.8)	(1.1)	(2.5)	(100.0)				
50～	14	37	52	44	56	52	48	40	49	95	2	64	42	34	27	2	4	14	16	6	698	11	709	205.25	173.93
(%)	(2.0)	(5.3)	(7.4)	(6.3)	(8.0)	(7.4)	(6.9)	(5.7)	(7.0)	(13.6)	(0.3)	(9.2)	(6.0)	(4.9)	(3.9)	(0.3)	(0.6)	(2.0)	(2.3)	(0.9)	(100.0)				
60～	9	27	21	27	22	41	30	24	19	67	2	35	33	21	14	6	6	8	8	7	427	7	434	220.85	198.69
(%)	(2.1)	(6.3)	(4.9)	(6.3)	(5.2)	(9.6)	(7.0)	(5.6)	(4.4)	(15.7)	(0.5)	(8.2)	(7.7)	(4.9)	(3.3)	(1.4)	(1.4)	(1.9)	(1.9)	(1.6)	(100.0)				
70～	4	7	6	19	17	16	13	13	14	31		23	19	5	3	1	1	2	3	8	205	8	213	238.61	323.47
(%)	(2.0)	(3.4)	(2.9)	(9.3)	(8.3)	(7.8)	(6.3)	(6.3)	(6.8)	(15.1)		(11.2)	(9.3)	(2.4)	(1.5)	(0.5)	(0.5)	(1.0)	(1.5)	(3.9)	(100.0)				
80～	4	4	4	8	9	7	7	5	3	19		10	11	10	3	1		2	3	5	115	2	117	255.21	246.49
(%)	(3.5)	(3.5)	(3.5)	(7.0)	(7.8)	(6.1)	(6.1)	(4.3)	(2.6)	(16.5)		(8.7)	(9.6)	(8.7)	(2.6)	(0.9)		(1.7)	(2.6)	(4.3)	(100.0)				
90～	1	1	4	4	4	6	1	4	2	7		4	3	4		3	1		1	1	51	1	52	223.76	189.21
(%)	(2.0)	(2.0)	(7.8)	(7.8)	(7.8)	(11.8)	(2.0)	(7.8)	(3.9)	(13.7)		(7.8)	(5.9)	(7.8)		(5.9)	(2.0)		(2.0)	(2.0)	(100.0)				
100～	2	4	3	4	6	3	5	1	2	9		2	3	2			1		1	1	49		49	180.20	162.87
(%)	(4.1)	(8.2)	(6.1)	(8.2)	(12.2)	(6.1)	(10.2)	(2.0)	(4.1)	(18.4)		(4.1)	(6.1)	(4.1)			(2.0)		(2.0)	(2.0)	(100.0)				
合計	92	192	199	219	235	251	223	208	204	468	7	312	229	149	95	30	32	55	58	61	3,319	73	3,392	214.74	219.64
(%)	(2.8)	(5.8)	(6.0)	(6.6)	(7.1)	(7.6)	(6.7)	(6.3)	(6.1)	(14.1)	(0.2)	(9.4)	(6.9)	(4.5)	(2.9)	(0.9)	(1.0)	(1.7)	(1.7)	(1.8)	(100.0)				
記載なし	31	89	69	96	74	65	82	77	72	186	3	119	102	51	44	13	22	18	47	49	1,309	303	1,612	247.04	250.76
(%)	(2.4)	(6.8)	(5.3)	(7.3)	(5.7)	(5.0)	(6.3)	(5.9)	(5.5)	(14.2)	(0.2)	(9.1)	(7.8)	(3.9)	(3.4)	(1.0)	(1.7)	(1.4)	(3.6)	(3.7)	(100.0)				
総計	123	281	268	315	309	316	305	285	276	654	10	431	331	200	139	43	54	73	105	110	4,628	376	5,004	223.88	229.31
(%)	(2.7)	(6.1)	(5.8)	(6.8)	(6.7)	(6.8)	(6.6)	(6.2)	(6.0)	(14.1)	(0.2)	(9.3)	(7.2)	(4.3)	(3.0)	(0.9)	(1.2)	(1.6)	(2.3)	(2.4)	(100.0)				
平均	49.29	48.47	48.56	51.72	51.83	51.63	50.35	49.16	49.28	51.65	49.57	49.93	51.57	51.95	49.79	54.67	50.63	48.75	52.48	54.90	50.67	47.75	50.61		
標準偏差	18.67	18.95	16.71	18.57	19.56	18.68	18.12	15.71	15.17	18.35	14.65	17.70	17.99	19.66	13.14	16.76	18.43	13.77	18.36	18.69	17.85	17.43	17.84		

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査