

表 3556 PTH値(pg/mL) 治療方法(新分類)別(透析患者全体) PTH測定法:whole-PTH

治療方法 (新分類)	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
施設血液透析	1,561	1,944	1,678	2,585	2,253	1,865	1,241	41	991	693	493	552	216	141	159	186	16,599	482	17,081	106.45	113.69
(%)	(9.4)	(11.7)	(10.1)	(15.6)	(13.6)	(11.2)	(7.5)	(0.2)	(6.0)	(4.2)	(3.0)	(3.3)	(1.3)	(0.8)	(1.0)	(1.1)	(100.0)				
Off-line HDF	42	48	33	48	45	36	20		23	17	7	11	1	7	3	7	348	5	353	106.05	114.20
(%)	(12.1)	(13.8)	(9.5)	(13.8)	(12.9)	(10.3)	(5.7)		(6.6)	(4.9)	(2.0)	(3.2)	(0.3)	(2.0)	(0.9)	(2.0)	(100.0)				
On-line HDF	217	268	248	402	358	301	212	4	199	127	91	80	46	22	24	40	2,639	26	2,665	113.42	112.61
(%)	(8.2)	(10.2)	(9.4)	(15.2)	(13.6)	(11.4)	(8.0)	(0.2)	(7.5)	(4.8)	(3.4)	(3.0)	(1.7)	(0.8)	(0.9)	(1.5)	(100.0)				
Push/Pull HDF												1					1		1	242.00	
(%)												(100.0)					(100.0)				
ハイオフィルトレーション	4	5	5	10	5	6	9			1	3	2		1		1	52	2	54	107.37	94.68
(%)	(7.7)	(9.6)	(9.6)	(19.2)	(9.6)	(11.5)	(17.3)			(1.9)	(5.8)	(3.8)		(1.9)		(1.9)	(100.0)				
血液濾過		1								1							2		2	114.00	115.97
(%)		(50.0)								(50.0)							(100.0)				
血液吸着透析 (リクセル)	26	16	16	24	16	14	9	2	5	2	4	3	1		4	1	143	1	144	91.77	103.33
(%)	(18.2)	(11.2)	(11.2)	(16.8)	(11.2)	(9.8)	(6.3)	(1.4)	(3.5)	(1.4)	(2.8)	(2.1)	(0.7)		(2.8)	(0.7)	(100.0)				
血液吸着透析 (その他)																					
(%)																					
在宅血液透析	4	4	8	7	2	5	3		2		3	2	1		1	1	43	1	44	113.42	112.66
(%)	(9.3)	(9.3)	(18.6)	(16.3)	(4.7)	(11.6)	(7.0)		(4.7)		(7.0)	(4.7)	(2.3)		(2.3)	(2.3)	(100.0)				
腹膜透析 (CAPD)	16	23	26	48	44	35	24		19	22	17	15	10	3	7	3	312	103	415	129.81	110.21
(%)	(5.1)	(7.4)	(8.3)	(15.4)	(14.1)	(11.2)	(7.7)		(6.1)	(7.1)	(5.4)	(4.8)	(3.2)	(1.0)	(2.2)	(1.0)	(100.0)				
腹膜透析 (APD)	18	13	19	21	18	21	17		18	9	7	7	5	4	3	7	187	28	215	143.99	163.61
(%)	(9.6)	(7.0)	(10.2)	(11.2)	(9.6)	(11.2)	(9.1)		(9.6)	(4.8)	(3.7)	(3.7)	(2.7)	(2.1)	(1.6)	(3.7)	(100.0)				
腹膜透析 (CCPD)	6	9	10	8	7	15	4	1	6	3	3	1	4	1	1	1	80	9	89	116.55	105.17
(%)	(7.5)	(11.3)	(12.5)	(10.0)	(8.8)	(18.8)	(5.0)	(1.3)	(7.5)	(3.8)	(3.8)	(1.3)	(5.0)	(1.3)	(1.3)	(1.3)	(100.0)				
合計	1,894	2,331	2,043	3,153	2,748	2,298	1,539	48	1,263	875	628	674	284	179	202	247	20,406	657	21,063	108.01	114.01
(%)	(9.3)	(11.4)	(10.0)	(15.5)	(13.5)	(11.3)	(7.5)	(0.2)	(6.2)	(4.3)	(3.1)	(3.3)	(1.4)	(0.9)	(1.0)	(1.2)	(100.0)				
記載なし (%)																					
総計	1,894	2,331	2,043	3,153	2,748	2,298	1,539	48	1,263	875	628	674	284	179	202	247	20,406	657	21,063	108.01	114.01
(%)	(9.3)	(11.4)	(10.0)	(15.5)	(13.5)	(11.3)	(7.5)	(0.2)	(6.2)	(4.3)	(3.1)	(3.3)	(1.4)	(0.9)	(1.0)	(1.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2014年調査