

表 3557 PTH値(pg/mL) 併用の状況別（透析患者全体）PTH測定法:whole-PTH

併用の状況	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	1,851	2,285	1,988	3,072	2,679	2,224	1,493	47	1,219	840	599	651	265	169	191	236	19,809	516	20,325	107.27	113.46
(%)	(9.3)	(11.5)	(10.0)	(15.5)	(13.5)	(11.2)	(7.5)	(0.2)	(6.2)	(4.2)	(3.0)	(3.3)	(1.3)	(0.9)	(1.0)	(1.2)	(100.0)				
PD単独	29	34	37	56	48	54	32	1	32	27	22	20	17	7	9	11	436	109	545	142.02	140.65
(%)	(6.7)	(7.8)	(8.5)	(12.8)	(11.0)	(12.4)	(7.3)	(0.2)	(7.3)	(6.2)	(5.0)	(4.6)	(3.9)	(1.6)	(2.1)	(2.5)	(100.0)				
HD(F)単独 PD力テ あり	1	1		2		2			1	1	2			1			11	1	12	140.09	107.79
(%)	(9.1)	(9.1)		(18.2)		(18.2)			(9.1)	(9.1)	(18.2)			(9.1)			(100.0)				
PD.HD併用週1 HD(F)	9	7	16	18	13	13	11		9	5	2	3	1	1	1		109	15	124	99.74	75.16
(%)	(8.3)	(6.4)	(14.7)	(16.5)	(11.9)	(11.9)	(10.1)		(8.3)	(4.6)	(1.8)	(2.8)	(0.9)	(0.9)	(0.9)		(100.0)				
PD.HD併用週2 HD(F)	2	4	2	4	7	4	2		2	2	3				1		33	16	49	109.36	88.57
(%)	(6.1)	(12.1)	(6.1)	(12.1)	(21.2)	(12.1)	(6.1)		(6.1)	(6.1)	(9.1)				(3.0)		(100.0)				
PD.HD併用週3 HD(F)	2			1			1										4		4	56.00	59.58
(%)	(50.0)			(25.0)			(25.0)										(100.0)				
PD.HD併用週4 HD(F)																					
(%)																					
PD.HD併用その他					1								1	1			3		3	261.33	163.22
(%)					(33.3)								(33.3)	(33.3)			(100.0)				
合計	1,894	2,331	2,043	3,153	2,748	2,297	1,539	48	1,263	875	628	674	284	179	202	247	20,405	657	21,062	108.01	114.02
(%)	(9.3)	(11.4)	(10.0)	(15.5)	(13.5)	(11.3)	(7.5)	(0.2)	(6.2)	(4.3)	(3.1)	(3.3)	(1.4)	(0.9)	(1.0)	(1.2)	(100.0)				
不明						1											1		1	113.00	
(%)						(100.0)											(100.0)				
記載なし																					
(%)																					
総計	1,894	2,331	2,043	3,153	2,748	2,298	1,539	48	1,263	875	628	674	284	179	202	247	20,406	657	21,063	108.01	114.01
(%)	(9.3)	(11.4)	(10.0)	(15.5)	(13.5)	(11.3)	(7.5)	(0.2)	(6.2)	(4.3)	(3.1)	(3.3)	(1.4)	(0.9)	(1.0)	(1.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査