

表 3561 PTH値(pg/mL) 原疾患別（透析患者全体）PTH測定法:whole-PTH

原疾患	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	581	695	648	980	899	714	503	17	427	297	224	250	111	68	72	113	6,599	245	6,844	115.57	128.16
(%)	(8.8)	(10.5)	(9.8)	(14.9)	(13.6)	(10.8)	(7.6)	(0.3)	(6.5)	(4.5)	(3.4)	(3.8)	(1.7)	(1.0)	(1.1)	(1.7)	(100.0)				
慢性腎盂腎炎	19	29	31	38	37	25	11	2	17	12	8	11	7	4	2	2	255	2	257	116.27	164.82
(%)	(7.5)	(11.4)	(12.2)	(14.9)	(14.5)	(9.8)	(4.3)	(0.8)	(6.7)	(4.7)	(3.1)	(4.3)	(2.7)	(1.6)	(0.8)	(0.8)	(100.0)				
急速進行性糸球体腎炎	32	29	28	31	27	15	11		10	4	2	3		1	2		195	6	201	76.61	70.11
(%)	(16.4)	(14.9)	(14.4)	(15.9)	(13.8)	(7.7)	(5.6)		(5.1)	(2.1)	(1.0)	(1.5)		(0.5)	(1.0)		(100.0)				
妊娠腎/妊娠中毒症	12	18	9	14	12	13	8		8	4	2	4	2		3	5	114	1	115	128.04	168.11
(%)	(10.5)	(15.8)	(7.9)	(12.3)	(10.5)	(11.4)	(7.0)		(7.0)	(3.5)	(1.8)	(3.5)	(1.8)		(2.6)	(4.4)	(100.0)				
その他の分類不能の腎炎	9	7	8	13	17	12	10		5	2	6	5	1	2	2	2	101	2	103	129.66	141.25
(%)	(8.9)	(6.9)	(7.9)	(12.9)	(16.8)	(11.9)	(9.9)		(5.0)	(2.0)	(5.9)	(5.0)	(1.0)	(2.0)	(2.0)	(2.0)	(100.0)				
多発性嚢胞腎	65	87	86	127	116	92	55	2	50	37	31	34	9	10	9	10	820	17	837	112.03	108.16
(%)	(7.9)	(10.6)	(10.5)	(15.5)	(14.1)	(11.2)	(6.7)	(0.2)	(6.1)	(4.5)	(3.8)	(4.1)	(1.1)	(1.2)	(1.1)	(1.2)	(100.0)				
腎硬化症	179	233	193	297	260	223	154	3	124	83	51	61	24	20	16	14	1,935	62	1,997	102.88	94.61
(%)	(9.3)	(12.0)	(10.0)	(15.3)	(13.4)	(11.5)	(8.0)	(0.2)	(6.4)	(4.3)	(2.6)	(3.2)	(1.2)	(1.0)	(0.8)	(0.7)	(100.0)				
悪性高血圧	12	22	13	29	30	16	15		16	8	3	10	2		3	6	185	7	192	126.67	138.06
(%)	(6.5)	(11.9)	(7.0)	(15.7)	(16.2)	(8.6)	(8.1)		(8.6)	(4.3)	(1.6)	(5.4)	(1.1)		(1.6)	(3.2)	(100.0)				
糖尿病性腎症	729	911	749	1,211	1,029	868	555	17	428	319	231	210	84	55	55	54	7,505	244	7,749	100.41	98.84
(%)	(9.7)	(12.1)	(10.0)	(16.1)	(13.7)	(11.6)	(7.4)	(0.2)	(5.7)	(4.3)	(3.1)	(2.8)	(1.1)	(0.7)	(0.7)	(0.7)	(100.0)				
SLE腎炎	13	18	20	21	10	21	12	1	10	5	8	7	3		3		152	7	159	108.13	90.77
(%)	(8.6)	(11.8)	(13.2)	(13.8)	(6.6)	(13.8)	(7.9)	(0.7)	(6.6)	(3.3)	(5.3)	(4.6)	(2.0)		(2.0)		(100.0)				
アミロイド腎	4	6	5	7	6	6	2	1		2						1	40	1	41	80.45	70.92
(%)	(10.0)	(15.0)	(12.5)	(17.5)	(15.0)	(15.0)	(5.0)	(2.5)		(5.0)						(2.5)	(100.0)				
痛風腎	7	6	3	11	9	13	5		5	1	7	3	1	1		1	73	1	74	118.85	94.14
(%)	(9.6)	(8.2)	(4.1)	(15.1)	(12.3)	(17.8)	(6.8)		(6.8)	(1.4)	(9.6)	(4.1)	(1.4)	(1.4)		(1.4)	(100.0)				
先天性代謝異常による腎不全	1	2	4	2	3	3	2		1	2	1		1	2			24		24	128.88	104.76
(%)	(4.2)	(8.3)	(16.7)	(8.3)	(12.5)	(12.5)	(8.3)		(4.2)	(8.3)	(4.2)		(4.2)	(8.3)			(100.0)				
腎・尿路結核	3	2	3	1	4		1		1	1		1	1		1		19	1	20	111.42	114.07
(%)	(15.8)	(10.5)	(15.8)	(5.3)	(21.1)		(5.3)		(5.3)	(5.3)		(5.3)	(5.3)		(5.3)		(100.0)				
腎・尿路結石	4	4	3	4	4	5	2		3	1	2	3					35	2	37	101.29	77.07
(%)	(11.4)	(11.4)	(8.6)	(11.4)	(11.4)	(14.3)	(5.7)		(8.6)	(2.9)	(5.7)	(8.6)					(100.0)				
腎・尿路腫瘍	3	6	3	5	9	9	5		1	6		1	2			1	51	1	52	114.49	94.16
(%)	(5.9)	(11.8)	(5.9)	(9.8)	(17.6)	(17.6)	(9.8)		(2.0)	(11.8)		(2.0)	(3.9)			(2.0)	(100.0)				
閉塞性尿路障害	5	5	9	8	3	7	4			2	3	2	2	1	1	1	53		53	118.77	124.36
(%)	(9.4)	(9.4)	(17.0)	(15.1)	(5.7)	(13.2)	(7.5)			(3.8)	(5.7)	(3.8)	(3.8)	(1.9)	(1.9)	(1.9)	(100.0)				
骨髄腫	5	3	7	6	1	1	2			1							26	1	27	56.88	44.02
(%)	(19.2)	(11.5)	(26.9)	(23.1)	(3.8)	(3.8)	(7.7)			(3.8)							(100.0)				
腎形成不全	7	6	5	7	6	3	3		3		2	1		1	1	2	47	2	49	133.94	219.70
(%)	(14.9)	(12.8)	(10.6)	(14.9)	(12.8)	(6.4)	(6.4)		(6.4)		(4.3)	(2.1)		(2.1)	(2.1)	(4.3)	(100.0)				
不明	144	176	153	238	198	185	129	4	109	63	34	42	24	9	21	23	1,552	47	1,599	110.64	123.33
(%)	(9.3)	(11.3)	(9.9)	(15.3)	(12.8)	(11.9)	(8.3)	(0.3)	(7.0)	(4.1)	(2.2)	(2.7)	(1.5)	(0.6)	(1.4)	(1.5)	(100.0)				
再導入	11	15	17	34	18	20	16		13	8	3	8	4	1	6	5	179		179	132.96	138.13
(%)	(6.1)	(8.4)	(9.5)	(19.0)	(10.1)	(11.2)	(8.9)		(7.3)	(4.5)	(1.7)	(4.5)	(2.2)	(0.6)	(3.4)	(2.8)	(100.0)				
その他	49	51	46	69	50	47	34	1	32	17	10	18	6	4	4	8	446	8	454	108.94	109.41
(%)	(11.0)	(11.4)	(10.3)	(15.5)	(11.2)	(10.5)	(7.6)	(0.2)	(7.2)	(3.8)	(2.2)	(4.0)	(1.3)	(0.9)	(0.9)	(1.8)	(100.0)				
合計	1,894	2,331	2,043	3,153	2,748	2,298	1,539	48	1,263	875	628	674	284	179	202	247	20,406	657	21,063	108.01	114.01
(%)	(9.3)	(11.4)	(10.0)	(15.5)	(13.5)	(11.3)	(7.5)	(0.2)	(6.2)	(4.3)	(3.1)	(3.3)	(1.4)	(0.9)	(1.0)	(1.2)	(100.0)				
記載なし																					
(%)																					
総計	1,894	2,331	2,043	3,153	2,748	2,298	1,539	48	1,263	875	628	674	284	179	202	247	20,406	657	21,063	108.01	114.01
(%)	(9.3)	(11.4)	(10.0)	(15.5)	(13.5)	(11.3)	(7.5)	(0.2)	(6.2)	(4.3)	(3.1)	(3.3)	(1.4)	(0.9)	(1.0)	(1.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査