

表 3564 PTH値(pg/mL) 透析前ヘモグロビン濃度(g/dL)別（透析患者全体）PTH測定法:whole-PTH

透析前ヘモグロビン濃度	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
5未満	3	1		1	2	1			1		1	1					11		11	98.36	89.16
(%)	(27.3)	(9.1)		(9.1)	(18.2)	(9.1)			(9.1)		(9.1)	(9.1)					(100.0)				
5～	1	2		1	1	1			2	1		1					10		10	112.40	84.85
(%)	(10.0)	(20.0)		(10.0)	(10.0)	(10.0)			(20.0)	(10.0)		(10.0)					(100.0)				
6～	11	5	2	6	5	1	4		4	1	4	6		1		1	51	9	60	120.18	128.35
(%)	(21.6)	(9.8)	(3.9)	(11.8)	(9.8)	(2.0)	(7.8)		(7.8)	(2.0)	(7.8)	(11.8)		(2.0)		(2.0)	(100.0)				
7～	32	16	26	33	33	23	17	2	10	7	8	8	4	4	4	4	231	14	245	113.95	129.52
(%)	(13.9)	(6.9)	(11.3)	(14.3)	(14.3)	(10.0)	(7.4)	(0.9)	(4.3)	(3.0)	(3.5)	(3.5)	(1.7)	(1.7)	(1.7)	(1.7)	(100.0)				
8～	104	118	82	114	98	78	61	2	54	32	31	25	12	8	6	19	844	38	882	112.31	160.03
(%)	(12.3)	(14.0)	(9.7)	(13.5)	(11.6)	(9.2)	(7.2)	(0.2)	(6.4)	(3.8)	(3.7)	(3.0)	(1.4)	(0.9)	(0.7)	(2.3)	(100.0)				
9～	282	344	280	484	394	314	204	5	169	120	92	102	35	27	47	49	2,948	52	3,000	111.56	123.47
(%)	(9.6)	(11.7)	(9.5)	(16.4)	(13.4)	(10.7)	(6.9)	(0.2)	(5.7)	(4.1)	(3.1)	(3.5)	(1.2)	(0.9)	(1.6)	(1.7)	(100.0)				
10～	605	760	692	1,046	928	804	557	15	431	270	194	200	99	58	53	68	6,780	82	6,862	105.31	104.26
(%)	(8.9)	(11.2)	(10.2)	(15.4)	(13.7)	(11.9)	(8.2)	(0.2)	(6.4)	(4.0)	(2.9)	(2.9)	(1.5)	(0.9)	(0.8)	(1.0)	(100.0)				
11～	543	695	645	976	845	727	444	16	387	276	191	206	80	52	55	64	6,202	82	6,284	106.44	108.09
(%)	(8.8)	(11.2)	(10.4)	(15.7)	(13.6)	(11.7)	(7.2)	(0.3)	(6.2)	(4.5)	(3.1)	(3.3)	(1.3)	(0.8)	(0.9)	(1.0)	(100.0)				
12～	216	273	225	360	306	248	191	7	136	125	72	85	37	20	21	18	2,340	38	2,378	106.63	98.06
(%)	(9.2)	(11.7)	(9.6)	(15.4)	(13.1)	(10.6)	(8.2)	(0.3)	(5.8)	(5.3)	(3.1)	(3.6)	(1.6)	(0.9)	(0.9)	(0.8)	(100.0)				
13～	58	65	45	77	70	52	33	1	34	22	13	24	11	2	9	10	526	7	533	115.52	147.53
(%)	(11.0)	(12.4)	(8.6)	(14.6)	(13.3)	(9.9)	(6.3)	(0.2)	(6.5)	(4.2)	(2.5)	(4.6)	(2.1)	(0.4)	(1.7)	(1.9)	(100.0)				
14～	20	33	18	28	24	18	11		17	6	9	10	2	5	4	3	208	5	213	118.76	139.65
(%)	(9.6)	(15.9)	(8.7)	(13.5)	(11.5)	(8.7)	(5.3)		(8.2)	(2.9)	(4.3)	(4.8)	(1.0)	(2.4)	(1.9)	(1.4)	(100.0)				
合計	1,875	2,312	2,015	3,126	2,706	2,267	1,522	48	1,245	860	615	668	280	177	199	236	20,151	327	20,478	107.56	112.63
(%)	(9.3)	(11.5)	(10.0)	(15.5)	(13.4)	(11.3)	(7.6)	(0.2)	(6.2)	(4.3)	(3.1)	(3.3)	(1.4)	(0.9)	(1.0)	(1.2)	(100.0)				
記載なし	19	19	28	27	42	31	17		18	15	13	6	4	2	3	11	255	330	585	143.45	191.50
(%)	(7.5)	(7.5)	(11.0)	(10.6)	(16.5)	(12.2)	(6.7)		(7.1)	(5.9)	(5.1)	(2.4)	(1.6)	(0.8)	(1.2)	(4.3)	(100.0)				
総計	1,894	2,331	2,043	3,153	2,748	2,298	1,539	48	1,263	875	628	674	284	179	202	247	20,406	657	21,063	108.01	114.01
(%)	(9.3)	(11.4)	(10.0)	(15.5)	(13.5)	(11.3)	(7.5)	(0.2)	(6.2)	(4.3)	(3.1)	(3.3)	(1.4)	(0.9)	(1.0)	(1.2)	(100.0)				
平均	10.76	10.84	10.83	10.85	10.84	10.85	10.83	10.79	10.85	10.92	10.79	10.89	10.89	10.81	10.79	10.63	10.83	10.45	10.83		
標準偏差	1.32	1.24	1.15	1.19	1.17	1.14	1.15	1.24	1.24	1.16	1.33	1.33	1.19	1.34	1.33	1.36	1.21	1.65	1.22		

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査