

表 3565 PTH値(pg/mL) 総コレステロール濃度(mg/dL)別 (透析患者全体) PTH測定法:whole-PTH

総コレステロール濃度	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
100未満	63	60	65	75	78	54	35	1	27	9	7	14	5	2	5	5	505	7	512	94.69	118.32
(%)	(12.5)	(11.9)	(12.9)	(14.9)	(15.4)	(10.7)	(6.9)	(0.2)	(5.3)	(1.8)	(1.4)	(2.8)	(1.0)	(0.4)	(1.0)	(1.0)	(100.0)				
100～	177	206	155	271	243	192	112	3	92	67	44	56	18	12	17	14	1,679	12	1,691	101.69	107.67
(%)	(10.5)	(12.3)	(9.2)	(16.1)	(14.5)	(11.4)	(6.7)	(0.2)	(5.5)	(4.0)	(2.6)	(3.3)	(1.1)	(0.7)	(1.0)	(0.8)	(100.0)				
120～	273	370	293	508	423	310	218	2	165	114	81	108	32	16	31	28	2,972	23	2,995	103.43	113.94
(%)	(9.2)	(12.4)	(9.9)	(17.1)	(14.2)	(10.4)	(7.3)	(0.1)	(5.6)	(3.8)	(2.7)	(3.6)	(1.1)	(0.5)	(1.0)	(0.9)	(100.0)				
140～	328	434	367	543	488	418	272	10	240	151	106	129	40	34	27	46	3,633	28	3,661	106.62	104.30
(%)	(9.0)	(11.9)	(10.1)	(14.9)	(13.4)	(11.5)	(7.5)	(0.3)	(6.6)	(4.2)	(2.9)	(3.6)	(1.1)	(0.9)	(0.7)	(1.3)	(100.0)				
160～	306	364	311	475	400	363	247	4	199	140	95	101	58	34	33	31	3,161	29	3,190	108.09	112.45
(%)	(9.7)	(11.5)	(9.8)	(15.0)	(12.7)	(11.5)	(7.8)	(0.1)	(6.3)	(4.4)	(3.0)	(3.2)	(1.8)	(1.1)	(1.0)	(1.0)	(100.0)				
180～	172	221	175	292	272	225	149	8	114	101	72	58	28	19	22	29	1,957	20	1,977	112.73	130.85
(%)	(8.8)	(11.3)	(8.9)	(14.9)	(13.9)	(11.5)	(7.6)	(0.4)	(5.8)	(5.2)	(3.7)	(3.0)	(1.4)	(1.0)	(1.1)	(1.5)	(100.0)				
200～	109	123	110	149	136	120	91	2	72	39	41	44	23	8	19	23	1,109	9	1,118	120.08	137.35
(%)	(9.8)	(11.1)	(9.9)	(13.4)	(12.3)	(10.8)	(8.2)	(0.2)	(6.5)	(3.5)	(3.7)	(4.0)	(2.1)	(0.7)	(1.7)	(2.1)	(100.0)				
220～	55	48	42	77	52	50	38	3	45	23	22	17	12	6	5	3	498	3	501	111.40	95.96
(%)	(11.0)	(9.6)	(8.4)	(15.5)	(10.4)	(10.0)	(7.6)	(0.6)	(9.0)	(4.6)	(4.4)	(3.4)	(2.4)	(1.2)	(1.0)	(0.6)	(100.0)				
240～	16	23	19	13	27	14	11		13	10	11	9	5	2	1	5	179	2	181	131.03	151.27
(%)	(8.9)	(12.8)	(10.6)	(7.3)	(15.1)	(7.8)	(6.1)		(7.3)	(5.6)	(6.1)	(5.0)	(2.8)	(1.1)	(0.6)	(2.8)	(100.0)				
260～	10	9	10	18	15	14	7	1	7	4	6	3		3	3	3	113	2	115	136.79	186.80
(%)	(8.8)	(8.0)	(8.8)	(15.9)	(13.3)	(12.4)	(6.2)	(0.9)	(6.2)	(3.5)	(5.3)	(2.7)		(2.7)	(2.7)	(2.7)	(100.0)				
合計	1,509	1,858	1,547	2,421	2,134	1,760	1,180	34	974	658	485	539	221	136	163	187	15,806	135	15,941	107.75	115.85
(%)	(9.5)	(11.8)	(9.8)	(15.3)	(13.5)	(11.1)	(7.5)	(0.2)	(6.2)	(4.2)	(3.1)	(3.4)	(1.4)	(0.9)	(1.0)	(1.2)	(100.0)				
記載なし	385	473	496	732	614	538	359	14	289	217	143	135	63	43	39	60	4,600	522	5,122	108.89	107.47
(%)	(8.4)	(10.3)	(10.8)	(15.9)	(13.3)	(11.7)	(7.8)	(0.3)	(6.3)	(4.7)	(3.1)	(2.9)	(1.4)	(0.9)	(0.8)	(1.3)	(100.0)				
総計	1,894	2,331	2,043	3,153	2,748	2,298	1,539	48	1,263	875	628	674	284	179	202	247	20,406	657	21,063	108.01	114.01
(%)	(9.3)	(11.4)	(10.0)	(15.5)	(13.5)	(11.3)	(7.5)	(0.2)	(6.2)	(4.3)	(3.1)	(3.3)	(1.4)	(0.9)	(1.0)	(1.2)	(100.0)				
平均	156.06	155.64	156.09	155.35	155.59	157.29	158.25	170.00	160.03	160.36	164.36	157.52	164.90	165.30	161.48	165.05	157.27	158.45	157.28		
標準偏差	36.32	34.59	35.85	34.58	35.06	35.10	34.95	39.95	36.16	34.62	37.64	35.61	34.96	36.53	38.28	37.27	35.44	42.45	35.50		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査