

表 3567 PTH値(pg/mL) Non-HDL-C濃度(mg/dL)別（透析患者全体）PTH測定法:whole-PTH

Non-HDL-C濃度	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
50未満	32	22	21	37	26	13	11	1	12	7	3	8	2	2	1	3	201	2	203	94.63	94.31
(%)	(15.9)	(10.9)	(10.4)	(18.4)	(12.9)	(6.5)	(5.5)	(0.5)	(6.0)	(3.5)	(1.5)	(4.0)	(1.0)	(1.0)	(0.5)	(1.5)	(100.0)				
50～	111	149	141	202	181	150	75		75	45	33	35	11	6	17	6	1,237	11	1,248	99.58	104.13
(%)	(9.0)	(12.0)	(11.4)	(16.3)	(14.6)	(12.1)	(6.1)		(6.1)	(3.6)	(2.7)	(2.8)	(0.9)	(0.5)	(1.4)	(0.5)	(100.0)				
70～	278	337	267	447	391	301	199	2	151	89	72	99	24	21	17	26	2,721	20	2,741	99.45	94.17
(%)	(10.2)	(12.4)	(9.8)	(16.4)	(14.4)	(11.1)	(7.3)	(0.1)	(5.5)	(3.3)	(2.6)	(3.6)	(0.9)	(0.8)	(0.6)	(1.0)	(100.0)				
90～	276	388	329	511	437	372	255	4	202	141	87	111	46	25	33	35	3,252	19	3,271	107.64	115.59
(%)	(8.5)	(11.9)	(10.1)	(15.7)	(13.4)	(11.4)	(7.8)	(0.1)	(6.2)	(4.3)	(2.7)	(3.4)	(1.4)	(0.8)	(1.0)	(1.1)	(100.0)				
110～	258	323	276	398	381	319	223	8	181	122	89	84	48	24	31	26	2,791	28	2,819	107.91	112.20
(%)	(9.2)	(11.6)	(9.9)	(14.3)	(13.7)	(11.4)	(8.0)	(0.3)	(6.5)	(4.4)	(3.2)	(3.0)	(1.7)	(0.9)	(1.1)	(0.9)	(100.0)				
130～	149	186	128	250	212	200	134	5	99	89	52	56	33	18	14	17	1,642	20	1,662	111.60	119.80
(%)	(9.1)	(11.3)	(7.8)	(15.2)	(12.9)	(12.2)	(8.2)	(0.3)	(6.0)	(5.4)	(3.2)	(3.4)	(2.0)	(1.1)	(0.9)	(1.0)	(100.0)				
150～	83	87	70	124	98	88	57	4	56	34	41	35	17	9	16	12	831	7	838	119.83	126.56
(%)	(10.0)	(10.5)	(8.4)	(14.9)	(11.8)	(10.6)	(6.9)	(0.5)	(6.7)	(4.1)	(4.9)	(4.2)	(2.0)	(1.1)	(1.9)	(1.4)	(100.0)				
170～	39	40	46	49	50	41	40		33	18	17	16	6	5	6	12	418	5	423	125.73	130.49
(%)	(9.3)	(9.6)	(11.0)	(11.7)	(12.0)	(9.8)	(9.6)		(7.9)	(4.3)	(4.1)	(3.8)	(1.4)	(1.2)	(1.4)	(2.9)	(100.0)				
190～	12	12	16	22	16	15	10	3	11	4	8	2	5		2	4	142	1	143	126.96	156.73
(%)	(8.5)	(8.5)	(11.3)	(15.5)	(11.3)	(10.6)	(7.0)	(2.1)	(7.7)	(2.8)	(5.6)	(1.4)	(3.5)		(1.4)	(2.8)	(100.0)				
210～	3	7	6	5	7	9	2		2	5	6	5		3	2	1	63	1	64	161.21	203.24
(%)	(4.8)	(11.1)	(9.5)	(7.9)	(11.1)	(14.3)	(3.2)		(3.2)	(7.9)	(9.5)	(7.9)		(4.8)	(3.2)	(1.6)	(100.0)				
合計	1,241	1,551	1,300	2,045	1,799	1,508	1,006	27	822	554	408	451	192	113	139	142	13,298	114	13,412	107.36	112.62
(%)	(9.3)	(11.7)	(9.8)	(15.4)	(13.5)	(11.3)	(7.6)	(0.2)	(6.2)	(4.2)	(3.1)	(3.4)	(1.4)	(0.8)	(1.0)	(1.1)	(100.0)				
記載なし	653	780	743	1,108	949	790	533	21	441	321	220	223	92	66	63	105	7,108	543	7,651	109.22	116.57
(%)	(9.2)	(11.0)	(10.5)	(15.6)	(13.4)	(11.1)	(7.5)	(0.3)	(6.2)	(4.5)	(3.1)	(3.1)	(1.3)	(0.9)	(0.9)	(1.5)	(100.0)				
総計	1,894	2,331	2,043	3,153	2,748	2,298	1,539	48	1,263	875	628	674	284	179	202	247	20,406	657	21,063	108.01	114.01
(%)	(9.3)	(11.4)	(10.0)	(15.5)	(13.5)	(11.3)	(7.5)	(0.2)	(6.2)	(4.3)	(3.1)	(3.3)	(1.4)	(0.9)	(1.0)	(1.2)	(100.0)				
平均	106.38	106.32	106.12	105.87	106.29	107.73	109.81	128.89	109.33	111.82	115.02	108.90	116.71	114.38	113.77	117.35	107.89	111.80	107.92		
標準偏差	33.39	31.79	33.09	32.41	32.20	33.22	32.01	36.39	32.93	33.04	37.13	33.80	31.97	36.62	36.47	37.39	33.04	35.13	33.05		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査