

表 3574 PTH値(pg/mL) 原疾患別（施設血液透析患者 透析回数週3回）PTH測定法:whole-PTH

原疾患	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	427	528	475	731	676	517	366	12	292	209	155	191	76	47	52	75	4,829	154	4,983	113.47	126.85
(%)	(8.8)	(10.9)	(9.8)	(15.1)	(14.0)	(10.7)	(7.6)	(0.2)	(6.0)	(4.3)	(3.2)	(4.0)	(1.6)	(1.0)	(1.1)	(1.6)	(100.0)				
慢性腎盂腎炎	15	24	23	27	24	19	7	2	10	10	6	6	3	4	1	1	182	2	184	113.25	184.46
(%)	(8.2)	(13.2)	(12.6)	(14.8)	(13.2)	(10.4)	(3.8)	(1.1)	(5.5)	(5.5)	(3.3)	(3.3)	(1.6)	(2.2)	(0.5)	(0.5)	(100.0)				
急速進行性糸球体腎炎	28	25	24	26	18	13	8		8	3	2	3					158	5	163	70.08	57.21
(%)	(17.7)	(15.8)	(15.2)	(16.5)	(11.4)	(8.2)	(5.1)		(5.1)	(1.9)	(1.3)	(1.9)					(100.0)				
妊娠腎/妊娠中毒症	8	10	7	10	8	8	6		5	1	2	3	2		2	2	74	1	75	122.91	164.13
(%)	(10.8)	(13.5)	(9.5)	(13.5)	(10.8)	(10.8)	(8.1)		(6.8)	(1.4)	(2.7)	(4.1)	(2.7)		(2.7)	(2.7)	(100.0)				
その他の分類不能の腎炎	6	5	7	9	14	8	9		4	2	2	5	1	2	2	1	77	1	78	132.91	149.10
(%)	(7.8)	(6.5)	(9.1)	(11.7)	(18.2)	(10.4)	(11.7)		(5.2)	(2.6)	(2.6)	(6.5)	(1.3)	(2.6)	(2.6)	(1.3)	(100.0)				
多発性嚢胞腎	56	73	71	100	88	73	44	1	38	25	18	31	7	8	8	9	650	8	658	110.82	110.30
(%)	(8.6)	(11.2)	(10.9)	(15.4)	(13.5)	(11.2)	(6.8)	(0.2)	(5.8)	(3.8)	(2.8)	(4.8)	(1.1)	(1.2)	(1.2)	(1.4)	(100.0)				
腎硬化症	139	190	162	240	219	192	128	2	103	66	39	54	19	15	14	13	1,595	42	1,637	103.56	93.35
(%)	(8.7)	(11.9)	(10.2)	(15.0)	(13.7)	(12.0)	(8.0)	(0.1)	(6.5)	(4.1)	(2.4)	(3.4)	(1.2)	(0.9)	(0.9)	(0.8)	(100.0)				
悪性高血圧	9	16	9	22	20	13	10		10	4	3	9	1		2	1	129	7	136	111.92	105.39
(%)	(7.0)	(12.4)	(7.0)	(17.1)	(15.5)	(10.1)	(7.8)		(7.8)	(3.1)	(2.3)	(7.0)	(0.8)		(1.6)	(0.8)	(100.0)				
糖尿病性腎症	610	739	614	987	833	692	444	14	331	260	188	160	67	45	39	38	6,061	178	6,239	98.47	91.38
(%)	(10.1)	(12.2)	(10.1)	(16.3)	(13.7)	(11.4)	(7.3)	(0.2)	(5.5)	(4.3)	(3.1)	(2.6)	(1.1)	(0.7)	(0.6)	(0.6)	(100.0)				
SLE腎炎	11	14	15	14	6	16	8	1	7	3	6	5	2		3		111	6	117	107.62	95.70
(%)	(9.9)	(12.6)	(13.5)	(12.6)	(5.4)	(14.4)	(7.2)	(0.9)	(6.3)	(2.7)	(5.4)	(4.5)	(1.8)		(2.7)		(100.0)				
アミロイド腎	2	4	4	4	5	6	1	1		1					1		29	1	30	87.03	76.16
(%)	(6.9)	(13.8)	(13.8)	(13.8)	(17.2)	(20.7)	(3.4)	(3.4)		(3.4)					(3.4)		(100.0)				
痛風腎	6	5	2	6	9	10	2		5		5	2	1	1		1	55	1	56	119.67	99.86
(%)	(10.9)	(9.1)	(3.6)	(10.9)	(16.4)	(18.2)	(3.6)		(9.1)		(9.1)	(3.6)	(1.8)	(1.8)		(1.8)	(100.0)				
先天性代謝異常による腎不全		2	2	1	3	2	1		1	2				2			16		16	133.06	107.95
(%)		(12.5)	(12.5)	(6.3)	(18.8)	(12.5)	(6.3)		(6.3)	(12.5)				(12.5)			(100.0)				
腎・尿路結核	3	2	3	1	4				1				1		1		16		16	95.94	116.68
(%)	(18.8)	(12.5)	(18.8)	(6.3)	(25.0)				(6.3)				(6.3)		(6.3)		(100.0)				
腎・尿路結石	4	3	1	4	3	4	1		2	1	2	2					27	2	29	99.30	77.22
(%)	(14.8)	(11.1)	(3.7)	(14.8)	(11.1)	(14.8)	(3.7)		(7.4)	(3.7)	(7.4)	(7.4)					(100.0)				
腎・尿路腫瘍	3	5	3	5	8	9	4		1	6		1	2			1	48		48	116.25	95.98
(%)	(6.3)	(10.4)	(6.3)	(10.4)	(16.7)	(18.8)	(8.3)		(2.1)	(12.5)		(2.1)	(4.2)			(2.1)	(100.0)				
閉塞性尿路障害	3	4	4	5	2	6	4			1	1	2	2		1	1	36		36	129.56	134.47
(%)	(8.3)	(11.1)	(11.1)	(13.9)	(5.6)	(16.7)	(11.1)			(2.8)	(2.8)	(5.6)	(5.6)		(2.8)	(2.8)	(100.0)				
骨髄腫	4	3	6	5	1	1	1			1							22	1	23	56.32	43.87
(%)	(18.2)	(13.6)	(27.3)	(22.7)	(4.5)	(4.5)	(4.5)			(4.5)							(100.0)				
腎形成不全	5	4	5	5	4	2	2		3		2	1		1	1	1	36		36	128.50	196.45
(%)	(13.9)	(11.1)	(13.9)	(13.9)	(11.1)	(5.6)	(5.6)		(8.3)		(5.6)	(2.8)		(2.8)	(2.8)	(2.8)	(100.0)				
不明	115	143	122	179	155	151	100	3	84	54	31	33	14	6	16	15	1,221	26	1,247	107.61	113.66
(%)	(9.4)	(11.7)	(10.0)	(14.7)	(12.7)	(12.4)	(8.2)	(0.2)	(6.9)	(4.4)	(2.5)	(2.7)	(1.1)	(0.5)	(1.3)	(1.2)	(100.0)				
再導入	7	10	13	25	11	15	12		9	5	2	4	3	1	6	3	126		126	131.73	129.66
(%)	(5.6)	(7.9)	(10.3)	(19.8)	(8.7)	(11.9)	(9.5)		(7.1)	(4.0)	(1.6)	(3.2)	(2.4)	(0.8)	(4.8)	(2.4)	(100.0)				
その他	38	39	41	53	37	30	29	1	20	12	8	14	4	3	3	4	336	3	339	103.25	100.89
(%)	(11.3)	(11.6)	(12.2)	(15.8)	(11.0)	(8.9)	(8.6)	(0.3)	(6.0)	(3.6)	(2.4)	(4.2)	(1.2)	(0.9)	(0.9)	(1.2)	(100.0)				
合計	1,499	1,848	1,613	2,459	2,148	1,787	1,187	37	934	666	472	526	205	135	152	166	15,834	438	16,272	105.70	109.24
(%)	(9.5)	(11.7)	(10.2)	(15.5)	(13.6)	(11.3)	(7.5)	(0.2)	(5.9)	(4.2)	(3.0)	(3.3)	(1.3)	(0.9)	(1.0)	(1.0)	(100.0)				
記載なし																					
(%)																					
総計	1,499	1,848	1,613	2,459	2,148	1,787	1,187	37	934	666	472	526	205	135	152	166	15,834	438	16,272	105.70	109.24
(%)	(9.5)	(11.7)	(10.2)	(15.5)	(13.6)	(11.3)	(7.5)	(0.2)	(5.9)	(4.2)	(3.0)	(3.3)	(1.3)	(0.9)	(1.0)	(1.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy
2014年調査