

表 3576 PTH値(pg/mL) 透析後体重(kg)別（施設血液透析患者 透析回数週3回）PTH測定法:whole-PTH

透析後体重	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
20未満	1		2	2	1												6		6	52.00	21.71
(%)	(16.7)		(33.3)	(33.3)	(16.7)												(100.0)				
20～	6	12	9	5	3	3	5		3	2	2	1		2		1	54	5	59	104.65	140.55
(%)	(11.1)	(22.2)	(16.7)	(9.3)	(5.6)	(5.6)	(9.3)		(5.6)	(3.7)	(3.7)	(1.9)		(3.7)		(1.9)	(100.0)				
30～	193	194	161	186	174	115	70	1	70	46	29	31	14	9	18	18	1,329	59	1,388	97.82	131.11
(%)	(14.5)	(14.6)	(12.1)	(14.0)	(13.1)	(8.7)	(5.3)	(0.1)	(5.3)	(3.5)	(2.2)	(2.3)	(1.1)	(0.7)	(1.4)	(1.4)	(100.0)				
40～	463	539	443	639	489	438	257	8	205	168	109	139	39	31	38	47	4,052	135	4,187	101.66	120.44
(%)	(11.4)	(13.3)	(10.9)	(15.8)	(12.1)	(10.8)	(6.3)	(0.2)	(5.1)	(4.1)	(2.7)	(3.4)	(1.0)	(0.8)	(0.9)	(1.2)	(100.0)				
50～	435	605	531	794	744	577	404	13	321	194	132	160	71	45	46	49	5,121	116	5,237	104.31	101.41
(%)	(8.5)	(11.8)	(10.4)	(15.5)	(11.3)	(7.9)	(6.3)	(0.3)	(6.3)	(3.8)	(2.6)	(3.1)	(1.4)	(0.9)	(0.9)	(1.0)	(100.0)				
60～	268	316	288	517	456	391	272	9	197	148	112	116	39	20	28	31	3,208	69	3,277	107.84	93.93
(%)	(8.4)	(9.9)	(9.0)	(16.1)	(14.2)	(12.2)	(8.5)	(0.3)	(6.1)	(4.6)	(3.5)	(3.6)	(1.2)	(0.6)	(0.9)	(1.0)	(100.0)				
70～	87	121	113	211	160	168	118	4	94	66	53	44	17	15	13	12	1,296	13	1,309	115.03	107.46
(%)	(6.7)	(9.3)	(8.7)	(16.3)	(12.3)	(13.0)	(9.1)	(0.3)	(7.3)	(5.1)	(4.1)	(3.4)	(1.3)	(1.2)	(1.0)	(0.9)	(100.0)				
80～	28	39	47	74	72	54	30	1	28	24	17	20	15	8	6	6	469	5	474	122.40	116.73
(%)	(6.0)	(8.3)	(10.0)	(15.8)	(15.4)	(11.5)	(6.4)	(0.2)	(6.0)	(5.1)	(3.6)	(4.3)	(3.2)	(1.7)	(1.3)	(1.3)	(100.0)				
90～	8	13	9	6	25	21	19	1	7	12	13	9	7	3	1	1	155	3	158	142.82	110.66
(%)	(5.2)	(8.4)	(5.8)	(3.9)	(16.1)	(13.5)	(12.3)	(0.6)	(4.5)	(7.7)	(8.4)	(5.8)	(4.5)	(1.9)	(0.6)	(0.6)	(100.0)				
100～	7	2	3	8	6	6	3		4	1	2	2	1	1		1	47		47	117.74	122.48
(%)	(14.9)	(4.3)	(6.4)	(17.0)	(12.8)	(12.8)	(6.4)		(8.5)	(2.1)	(4.3)	(4.3)	(2.1)	(2.1)		(2.1)	(100.0)				
110～	1			5	5	1	1		1	2	1	1	2		1		21		21	151.00	111.98
(%)	(4.8)			(23.8)	(23.8)	(4.8)	(4.8)		(4.8)	(9.5)	(4.8)	(4.8)	(9.5)		(4.8)		(100.0)				
120～		2	1	3	4	2			1			2			1		16		16	129.44	124.80
(%)		(12.5)	(6.3)	(18.8)	(25.0)	(12.5)			(6.3)			(12.5)			(6.3)		(100.0)				
合計	1,497	1,843	1,607	2,450	2,139	1,776	1,179	37	931	663	470	525	205	134	152	166	15,774	405	16,179	105.70	109.37
(%)	(9.5)	(11.7)	(10.2)	(15.5)	(13.6)	(11.3)	(7.5)	(0.2)	(5.9)	(4.2)	(3.0)	(3.3)	(1.3)	(0.8)	(1.0)	(1.1)	(100.0)				
記載なし	2	5	6	9	9	11	8		3	3	2	1		1			60	33	93	104.02	65.83
(%)	(3.3)	(8.3)	(10.0)	(15.0)	(15.0)	(18.3)	(13.3)		(5.0)	(5.0)	(3.3)	(1.7)		(1.7)			(100.0)				
総計	1,499	1,848	1,613	2,459	2,148	1,787	1,187	37	934	666	472	526	205	135	152	166	15,834	438	16,272	105.70	109.24
(%)	(9.5)	(11.7)	(10.2)	(15.5)	(13.6)	(11.3)	(7.5)	(0.2)	(5.9)	(4.2)	(3.0)	(3.3)	(1.3)	(0.9)	(1.0)	(1.0)	(100.0)				
平均	53.10	53.83	54.40	55.86	56.47	56.84	57.33	58.19	56.89	57.30	58.48	57.13	59.68	57.33	55.98	54.95	55.82	51.51	55.71		
標準偏差	12.44	12.44	12.41	12.62	13.21	12.76	12.64	12.36	12.95	13.57	13.75	13.91	15.15	14.68	15.23	12.90	12.97	11.39	12.95		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査