

表 3577 PTH値(pg/mL) 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回）PTH測定法:whole-PTH

透析前ヘモグロビン濃度	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
5未満	2	1		1	2	1			1		1	1					10		10	106.50	89.58
(%)	(20.0)	(10.0)		(10.0)	(20.0)	(10.0)			(10.0)		(10.0)	(10.0)					(100.0)				
5～	1	1		1		1			2	1		1					8		8	127.00	88.66
(%)	(12.5)	(12.5)		(12.5)		(12.5)			(25.0)	(12.5)		(12.5)					(100.0)				
6～	8	3	2	4	3	1	4		4	1	3	5		1		1	40	8	48	131.32	136.16
(%)	(20.0)	(7.5)	(5.0)	(10.0)	(7.5)	(2.5)	(10.0)		(10.0)	(2.5)	(7.5)	(12.5)		(2.5)		(2.5)	(100.0)				
7～	26	14	21	29	26	23	15	1	7	7	8	6	4	4	2	3	196	11	207	112.00	121.75
(%)	(13.3)	(7.1)	(10.7)	(14.8)	(13.3)	(11.7)	(7.7)	(0.5)	(3.6)	(3.6)	(4.1)	(3.1)	(2.0)	(2.0)	(1.0)	(1.5)	(100.0)				
8～	90	96	65	85	75	63	47	1	36	29	25	15	8	5	5	10	655	34	689	104.88	154.19
(%)	(13.7)	(14.7)	(9.9)	(13.0)	(11.5)	(9.6)	(7.2)	(0.2)	(5.5)	(4.4)	(3.8)	(2.3)	(1.2)	(0.8)	(0.8)	(1.5)	(100.0)				
9～	221	276	219	382	318	251	163	4	121	84	66	82	27	26	36	32	2,308	44	2,352	108.71	115.11
(%)	(9.6)	(12.0)	(9.5)	(16.6)	(13.8)	(10.9)	(7.1)	(0.2)	(5.2)	(3.6)	(2.9)	(3.6)	(1.2)	(1.1)	(1.6)	(1.4)	(100.0)				
10～	482	611	555	813	732	629	436	11	329	214	148	151	74	40	40	46	5,311	66	5,377	102.92	95.54
(%)	(9.1)	(11.5)	(10.5)	(15.3)	(13.8)	(11.8)	(8.2)	(0.2)	(6.2)	(4.0)	(2.8)	(2.8)	(1.4)	(0.8)	(0.8)	(0.9)	(100.0)				
11～	426	535	505	773	665	548	333	13	279	208	147	163	55	38	42	43	4,773	67	4,840	104.34	104.62
(%)	(8.9)	(11.2)	(10.6)	(16.2)	(13.9)	(11.5)	(7.0)	(0.3)	(5.8)	(4.4)	(3.1)	(3.4)	(1.2)	(0.8)	(0.9)	(0.9)	(100.0)				
12～	173	225	172	274	230	199	144	6	100	89	53	67	25	13	17	13	1,800	35	1,835	104.46	96.41
(%)	(9.6)	(12.5)	(9.6)	(15.2)	(12.8)	(11.1)	(8.0)	(0.3)	(5.6)	(4.9)	(2.9)	(3.7)	(1.4)	(0.7)	(0.9)	(0.7)	(100.0)				
13～	43	47	37	56	52	34	25	1	28	20	5	21	7	2	5	5	388	4	392	112.98	152.66
(%)	(11.1)	(12.1)	(9.5)	(14.4)	(13.4)	(8.8)	(6.4)	(0.3)	(7.2)	(5.2)	(1.3)	(5.4)	(1.8)	(0.5)	(1.3)	(1.3)	(100.0)				
14～	13	21	13	21	15	9	8		11	3	6	9	1	4	2	2	138	5	143	120.99	148.32
(%)	(9.4)	(15.2)	(9.4)	(15.2)	(10.9)	(6.5)	(5.8)		(8.0)	(2.2)	(4.3)	(6.5)	(0.7)	(2.9)	(1.4)	(1.4)	(100.0)				
合計	1,485	1,830	1,589	2,439	2,118	1,759	1,175	37	918	656	462	521	201	133	149	155	15,627	274	15,901	105.08	107.17
(%)	(9.5)	(11.7)	(10.2)	(15.6)	(13.6)	(11.3)	(7.5)	(0.2)	(5.9)	(4.2)	(3.0)	(3.3)	(1.3)	(0.9)	(1.0)	(1.0)	(100.0)				
記載なし	14	18	24	20	30	28	12		16	10	10	5	4	2	3	11	207	164	371	152.33	209.24
(%)	(6.8)	(8.7)	(11.6)	(9.7)	(14.5)	(13.5)	(5.8)		(7.7)	(4.8)	(4.8)	(2.4)	(1.9)	(1.0)	(1.4)	(5.3)	(100.0)				
総計	1,499	1,848	1,613	2,459	2,148	1,787	1,187	37	934	666	472	526	205	135	152	166	15,834	438	16,272	105.70	109.24
(%)	(9.5)	(11.7)	(10.2)	(15.5)	(13.6)	(11.3)	(7.5)	(0.2)	(5.9)	(4.2)	(3.0)	(3.3)	(1.3)	(0.9)	(1.0)	(1.0)	(100.0)				
平均	10.74	10.81	10.82	10.84	10.82	10.81	10.80	10.93	10.85	10.88	10.73	10.91	10.82	10.73	10.77	10.64	10.81	10.43	10.81		
標準偏差	1.31	1.21	1.16	1.18	1.15	1.13	1.16	1.13	1.23	1.18	1.33	1.36	1.20	1.41	1.25	1.34	1.20	1.69	1.21		

数値下のかつこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査