

表 3578 PTH値(pg/mL) 総コレステロール濃度(mg/dL)別（施設血液透析患者 透析回数週3回）PTH測定法:whole-PTH

総コレステロール濃度	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
100未満	54	47	54	57	60	39	25		22	7	7	11	4	2	3	4	396	5	401	94.19	126.14
(%)	(13.6)	(11.9)	(13.6)	(14.4)	(15.2)	(9.8)	(6.3)		(5.6)	(1.8)	(1.8)	(2.8)	(1.0)	(0.5)	(0.8)	(1.0)	(100.0)				
100～	144	162	126	209	191	147	90	2	69	55	37	44	13	8	16	9	1,322	12	1,334	100.89	107.32
(%)	(10.9)	(12.3)	(9.5)	(15.8)	(14.4)	(11.1)	(6.8)	(0.2)	(5.2)	(4.2)	(2.8)	(3.3)	(1.0)	(0.6)	(1.2)	(0.7)	(100.0)				
120～	230	297	232	404	337	238	166	2	116	85	65	83	22	10	20	21	2,328	20	2,348	99.74	103.59
(%)	(9.9)	(12.8)	(10.0)	(17.4)	(14.5)	(10.2)	(7.1)	(0.1)	(5.0)	(3.7)	(2.8)	(3.6)	(0.9)	(0.4)	(0.9)	(0.9)	(100.0)				
140～	253	345	298	431	374	325	211	7	178	114	76	100	32	24	22	33	2,823	22	2,845	104.85	102.04
(%)	(9.0)	(12.2)	(10.6)	(15.3)	(13.2)	(11.5)	(7.5)	(0.2)	(6.3)	(4.0)	(2.7)	(3.5)	(1.1)	(0.9)	(0.8)	(1.2)	(100.0)				
160～	227	283	234	363	301	289	189	3	148	109	67	82	39	26	25	19	2,404	18	2,422	107.16	113.47
(%)	(9.4)	(11.8)	(9.7)	(15.1)	(12.5)	(12.0)	(7.9)	(0.1)	(6.2)	(4.5)	(2.8)	(3.4)	(1.6)	(1.1)	(1.0)	(0.8)	(100.0)				
180～	141	167	129	216	223	159	116	7	81	81	44	49	18	15	14	17	1,477	14	1,491	108.43	114.90
(%)	(9.5)	(11.3)	(8.7)	(14.6)	(15.1)	(10.8)	(7.9)	(0.5)	(5.5)	(5.5)	(3.0)	(3.3)	(1.2)	(1.0)	(0.9)	(1.2)	(100.0)				
200～	75	94	81	111	101	85	65		50	28	24	33	19	4	13	13	796	7	803	116.37	135.26
(%)	(9.4)	(11.8)	(10.2)	(13.9)	(12.7)	(10.7)	(8.2)		(6.3)	(3.5)	(3.0)	(4.1)	(2.4)	(0.5)	(1.6)	(1.6)	(100.0)				
220～	42	37	31	60	36	36	25	2	33	14	17	12	9	5	3	1	363	2	365	107.47	91.67
(%)	(11.6)	(10.2)	(8.5)	(16.5)	(9.9)	(9.9)	(6.9)	(0.6)	(9.1)	(3.9)	(4.7)	(3.3)	(2.5)	(1.4)	(0.8)	(0.3)	(100.0)				
240～	12	17	10	9	18	11	8		10	6	9	8	5	2	1	2	128		128	136.24	158.01
(%)	(9.4)	(13.3)	(7.8)	(7.0)	(14.1)	(8.6)	(6.3)		(7.8)	(4.7)	(7.0)	(6.3)	(3.9)	(1.6)	(0.8)	(1.6)	(100.0)				
260～	5	6	7	14	9	11	4	1	7	3	5	3		3	2	1	81	1	82	126.17	107.58
(%)	(6.2)	(7.4)	(8.6)	(17.3)	(11.1)	(13.6)	(4.9)	(1.2)	(8.6)	(3.7)	(6.2)	(3.7)		(3.7)	(2.5)	(1.2)	(100.0)				
合計	1,183	1,455	1,202	1,874	1,650	1,340	899	24	714	502	351	425	161	99	119	120	12,118	101	12,219	105.29	110.66
(%)	(9.8)	(12.0)	(9.9)	(15.5)	(13.6)	(11.1)	(7.4)	(0.2)	(5.9)	(4.1)	(2.9)	(3.5)	(1.3)	(0.8)	(1.0)	(1.0)	(100.0)				
記載なし	316	393	411	585	498	447	288	13	220	164	121	101	44	36	33	46	3,716	337	4,053	107.01	104.47
(%)	(8.5)	(10.6)	(11.1)	(15.7)	(13.4)	(12.0)	(7.8)	(0.3)	(5.9)	(4.4)	(3.3)	(2.7)	(1.2)	(1.0)	(0.9)	(1.2)	(100.0)				
総計	1,499	1,848	1,613	2,459	2,148	1,787	1,187	37	934	666	472	526	205	135	152	166	15,834	438	16,272	105.70	109.24
(%)	(9.5)	(11.7)	(10.2)	(15.5)	(13.6)	(11.3)	(7.5)	(0.2)	(5.9)	(4.2)	(3.0)	(3.3)	(1.3)	(0.9)	(1.0)	(1.0)	(100.0)				
平均	154.45	154.95	154.36	154.82	154.92	156.69	157.39	172.17	160.02	159.43	161.99	158.08	165.62	167.09	159.50	160.05	156.36	153.62	156.34		
標準偏差	35.93	34.14	34.61	34.49	34.36	34.55	34.18	38.07	36.84	33.89	39.28	35.83	36.34	38.34	38.22	34.91	35.06	41.13	35.12		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査