

表 3587 PTH値(pg/mL) 原疾患別（血液透析濾過患者 透析回数週3回）PTH測定法whole-PTH

原疾患	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	100	112	109	170	155	139	93	2	94	62	47	42	18	15	12	23	1,193	17	1,210	118.19	110.22
(%)	(8.4)	(9.4)	(9.1)	(14.2)	(13.0)	(11.7)	(7.8)	(0.2)	(7.9)	(5.2)	(3.9)	(3.5)	(1.5)	(1.3)	(1.0)	(1.9)	(100.0)				
慢性腎盂腎炎	3		4	5	6		2		5	1	2	2	1			1	32		32	128.22	104.04
(%)	(9.4)		(12.5)	(15.6)	(18.8)		(6.3)		(15.6)	(3.1)	(6.3)	(6.3)	(3.1)			(3.1)	(100.0)				
急速進行性糸球体腎炎	3	4	3	1	4	2	1		2	1				1	1		23		23	103.26	112.86
(%)	(13.0)	(17.4)	(13.0)	(4.3)	(17.4)	(8.7)	(4.3)		(8.7)	(4.3)				(4.3)	(4.3)		(100.0)				
妊娠腎/妊娠中毒症	3	5	1	2	3	2	2		2	3		1			1	1	26		26	124.19	129.33
(%)	(11.5)	(19.2)	(3.8)	(7.7)	(11.5)	(7.7)	(7.7)		(7.7)	(11.5)		(3.8)			(3.8)	(3.8)	(100.0)				
その他の分類不能の腎炎	1		1	1		2						4					9		9	138.33	89.70
(%)	(11.1)		(11.1)	(11.1)		(22.2)						(44.4)					(100.0)				
多発性嚢胞腎	5	10	13	19	19	15	10	1	9	10	9	2	1	2	1	1	127	1	128	120.82	106.69
(%)	(3.9)	(7.9)	(10.2)	(15.0)	(15.0)	(11.8)	(7.9)	(0.8)	(7.1)	(7.9)	(7.1)	(1.6)	(0.8)	(1.6)	(0.8)	(0.8)	(100.0)				
腎硬化症	21	24	21	40	22	19	16		10	8	6	3	3	3	1		197	3	200	93.05	78.37
(%)	(10.7)	(12.2)	(10.7)	(20.3)	(11.2)	(9.6)	(8.1)		(5.1)	(4.1)	(3.0)	(1.5)	(1.5)	(1.5)	(0.5)		(100.0)				
悪性高血圧	2	1	2	2	7	2	3		1	2						2	24		24	138.92	161.10
(%)	(8.3)	(4.2)	(8.3)	(8.3)	(29.2)	(8.3)	(12.5)		(4.2)	(8.3)						(8.3)	(100.0)				
糖尿病性腎症	78	106	89	137	126	106	68		61	36	25	24	10	5	5	7	883	3	886	99.64	85.79
(%)	(8.8)	(12.0)	(10.1)	(15.5)	(14.3)	(12.0)	(7.7)		(6.9)	(4.1)	(2.8)	(2.7)	(1.1)	(0.6)	(0.6)	(0.8)	(100.0)				
SLE腎炎	1	3	4	5	2	3	4		2	1	1	2	1				29	1	30	112.34	82.62
(%)	(3.4)	(10.3)	(13.8)	(17.2)	(6.9)	(10.3)	(13.8)		(6.9)	(3.4)	(3.4)	(6.9)	(3.4)				(100.0)				
アミロイド腎	1	2		3			1										7		7	53.71	40.94
(%)	(14.3)	(28.6)		(42.9)			(14.3)										(100.0)				
痛風腎	1	1		3		2	1				1	1					10		10	112.30	85.90
(%)	(10.0)	(10.0)		(30.0)		(20.0)	(10.0)				(10.0)	(10.0)					(100.0)				
先天性代謝異常による腎不全	1		1	1							1						4		4	88.25	94.81
(%)	(25.0)		(25.0)	(25.0)							(25.0)						(100.0)				
腎・尿路結核							1			1			1				3	1	4	194.00	53.51
(%)							(33.3)			(33.3)			(33.3)				(100.0)				
腎・尿路結石			2		1	1	1						1				6		6	114.50	86.48
(%)			(33.3)		(16.7)	(16.7)	(16.7)						(16.7)				(100.0)				
腎・尿路腫瘍		1			1		1										3		3	86.33	62.58
(%)		(33.3)			(33.3)		(33.3)										(100.0)				
閉塞性尿路障害	1	1	2		1	1				1	1			1			9		9	121.89	120.11
(%)	(11.1)	(11.1)	(22.2)		(11.1)	(11.1)				(11.1)	(11.1)			(11.1)			(100.0)				
骨髄腫	1			1													2		2	38.50	44.55
(%)	(50.0)			(50.0)													(100.0)				
腎形成不全	1	2		1	1	1											6		6	57.67	39.01
(%)	(16.7)	(33.3)		(16.7)	(16.7)	(16.7)											(100.0)				
不明	19	20	15	37	30	21	23	1	17	5	1	3	6	1	2	6	207	4	211	122.48	166.01
(%)	(9.2)	(9.7)	(7.2)	(17.9)	(14.5)	(10.1)	(11.1)	(0.5)	(8.2)	(2.4)	(0.5)	(1.4)	(2.9)	(0.5)	(1.0)	(2.9)	(100.0)				
再導入	1	3	3	5	4	2	3		4	3		3				1	32		32	137.41	140.53
(%)	(3.1)	(9.4)	(9.4)	(15.6)	(12.5)	(6.3)	(9.4)		(12.5)	(9.4)		(9.4)				(3.1)	(100.0)				
その他	6	6	4	11	9	13	4		8	3		3	2	1	1	3	74	1	75	135.38	139.96
(%)	(8.1)	(8.1)	(5.4)	(14.9)	(12.2)	(17.6)	(5.4)		(10.8)	(4.1)		(4.1)	(2.7)	(1.4)	(1.4)	(4.1)	(100.0)				
合計	249	301	274	444	391	331	234	4	215	137	98	88	42	29	24	45	2,906	31	2,937	111.79	108.12
(%)	(8.6)	(10.4)	(9.4)	(15.3)	(13.5)	(11.4)	(8.1)	(0.1)	(7.4)	(4.7)	(3.4)	(3.0)	(1.4)	(1.0)	(0.8)	(1.5)	(100.0)				
記載なし																					
(%)																					
総計	249	301	274	444	391	331	234	4	215	137	98	88	42	29	24	45	2,906	31	2,937	111.79	108.12
(%)	(8.6)	(10.4)	(9.4)	(15.3)	(13.5)	(11.4)	(8.1)	(0.1)	(7.4)	(4.7)	(3.4)	(3.0)	(1.4)	(1.0)	(0.8)	(1.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査