

表 3600 PTH値(pg/mL) 原疾患別（腹膜透析患者）PTH測定法:whole-PTH

原疾患	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	12	20	22	30	29	25	18	1	13	17	9	5	10	4	3	2	220	62	282	127.31	124.15
(%)	(5.5)	(9.1)	(10.0)	(13.6)	(13.2)	(11.4)	(8.2)	(0.5)	(5.9)	(7.7)	(4.1)	(2.3)	(4.5)	(1.8)	(1.4)	(0.9)	(100.0)				
慢性腎盂腎炎			2		1	1			1			1					6		6	120.33	95.16
(%)			(33.3)		(16.7)	(16.7)			(16.7)			(16.7)					(100.0)				
急速進行性糸球体腎炎	1		1	1	1		1										5		5	68.80	47.74
(%)	(20.0)		(20.0)	(20.0)	(20.0)		(20.0)										(100.0)				
妊娠腎/妊娠中毒症		1			1	2			1							1	6		6	234.67	324.53
(%)		(16.7)			(16.7)	(33.3)			(16.7)							(16.7)	(100.0)				
その他の分類不能の腎炎				3	1				1							1	6		6	166.00	190.21
(%)				(50.0)	(16.7)				(16.7)							(16.7)	(100.0)				
多発性嚢胞腎	2			1	1	1			2	1			1				9	4	13	128.56	99.99
(%)	(22.2)			(11.1)	(11.1)	(11.1)			(22.2)	(11.1)			(11.1)				(100.0)				
腎硬化症	5	4	3	7	8	6	5		4	4	4	3	2	2			57	15	72	126.51	91.85
(%)	(8.8)	(7.0)	(5.3)	(12.3)	(14.0)	(10.5)	(8.8)		(7.0)	(7.0)	(7.0)	(5.3)	(3.5)	(3.5)			(100.0)				
悪性高血圧	1			1			1		2				1			1	7		7	195.00	167.15
(%)	(14.3)			(14.3)			(14.3)		(28.6)				(14.3)			(14.3)	(100.0)				
糖尿病性腎症	12	17	18	23	19	24	12		13	10	9	7	3	1	6	3	177	45	222	127.33	117.51
(%)	(6.8)	(9.6)	(10.2)	(13.0)	(10.7)	(13.6)	(6.8)		(7.3)	(5.6)	(5.1)	(4.0)	(1.7)	(0.6)	(3.4)	(1.7)	(100.0)				
SLE腎炎			1		2	2			1								6		6	105.50	44.24
(%)			(16.7)		(33.3)	(33.3)			(16.7)								(100.0)				
アミロイド腎			1														1		1	35.00	
(%)			(100.0)														(100.0)				
痛風腎			1	1			2										4		4	96.00	53.51
(%)			(25.0)	(25.0)			(50.0)										(100.0)				
先天性代謝異常による腎不全			1			1	1										3		3	96.67	43.25
(%)			(33.3)			(33.3)	(33.3)										(100.0)				
腎・尿路結核																					
(%)																					
腎・尿路結石																					
(%)																					
腎・尿路腫瘍																					
(%)																					
閉塞性尿路障害			1	1							1						3		3	110.33	99.65
(%)			(33.3)	(33.3)							(33.3)						(100.0)				
骨髄腫							1										1		1	127.00	
(%)							(100.0)										(100.0)				
腎形成不全					1		1									1	3	2	5	418.00	530.69
(%)					(33.3)		(33.3)									(33.3)	(100.0)				
不明	3	2	4	7	4	7	2		4	2	1	5	2	1	2	1	47	8	55	156.45	147.32
(%)	(6.4)	(4.3)	(8.5)	(14.9)	(8.5)	(14.9)	(4.3)		(8.5)	(4.3)	(2.1)	(10.6)	(4.3)	(2.1)	(4.3)	(2.1)	(100.0)				
再導入	1										1	1					3		3	167.00	134.73
(%)	(33.3)										(33.3)	(33.3)					(100.0)				
その他	3	1		2	1	2	1		1		2	1				1	15	4	19	139.60	150.73
(%)	(20.0)	(6.7)		(13.3)	(6.7)	(13.3)	(6.7)		(6.7)		(13.3)	(6.7)				(6.7)	(100.0)				
合計	40	45	55	77	69	71	45	1	43	34	27	23	19	8	11	11	579	140	719	132.56	129.39
(%)	(6.9)	(7.8)	(9.5)	(13.3)	(11.9)	(12.3)	(7.8)	(0.2)	(7.4)	(5.9)	(4.7)	(4.0)	(3.3)	(1.4)	(1.9)	(1.9)	(100.0)				
記載なし																					
(%)																					
総計	40	45	55	77	69	71	45	1	43	34	27	23	19	8	11	11	579	140	719	132.56	129.39
(%)	(6.9)	(7.8)	(9.5)	(13.3)	(11.9)	(12.3)	(7.8)	(0.2)	(7.4)	(5.9)	(4.7)	(4.0)	(3.3)	(1.4)	(1.9)	(1.9)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査