

表 3612 透析前ヘモグロビン濃度(g/dL) 併用の状況別（透析患者全体）

併用の状況	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	296	415	1,130	4,302	14,186	45,033	92,126	76,377	28,027	7,057	2,588	271,537	29,257	300,794	10.71	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.2)	(16.6)	(33.9)	(28.1)	(10.3)	(2.6)	(1.0)	(100.0)				
PD単独	6	12	23	94	287	707	1,330	1,315	596	187	60	4,617	2,571	7,188	10.79	1.43
(%)	(0.1)	(0.3)	(0.5)	(2.0)	(6.2)	(15.3)	(28.8)	(28.5)	(12.9)	(4.1)	(1.3)	(100.0)				
HD(F)単独.PDカテあり			2	7	18	41	62	59	17	8	4	218	51	269	10.57	1.47
(%)			(0.9)	(3.2)	(8.3)	(18.8)	(28.4)	(27.1)	(7.8)	(3.7)	(1.8)	(100.0)				
PD.HD併用週1 HD(F)	1	1	5	26	50	180	343	338	172	43	17	1,176	368	1,544	10.87	1.37
(%)	(0.1)	(0.1)	(0.4)	(2.2)	(4.3)	(15.3)	(29.2)	(28.7)	(14.6)	(3.7)	(1.4)	(100.0)				
PD.HD併用週2 HD(F)				1	5	13	25	51	29	9	2	135	42	177	11.32	1.27
(%)				(0.7)	(3.7)	(9.6)	(18.5)	(37.8)	(21.5)	(6.7)	(1.5)	(100.0)				
PD.HD併用週3 HD(F)				1	2	6	8	13	2	2		34	6	40	10.71	1.33
(%)				(2.9)	(5.9)	(17.6)	(23.5)	(38.2)	(5.9)	(5.9)		(100.0)				
PD.HD併用週4 HD(F)													1	1		
(%)																
PD.HD併用その他		1		3	6	6	14	17	9	1	2	59	14	73	10.63	1.75
(%)		(1.7)		(5.1)	(10.2)	(10.2)	(23.7)	(28.8)	(15.3)	(1.7)	(3.4)	(100.0)				
合計	303	429	1,160	4,434	14,554	45,986	93,908	78,170	28,852	7,307	2,673	277,776	32,310	310,086	10.71	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.2)	(16.6)	(33.8)	(28.1)	(10.4)	(2.6)	(1.0)	(100.0)				
不明				1		3	4	3	2	1		14	8	22	10.68	1.49
(%)				(7.1)		(21.4)	(28.6)	(21.4)	(14.3)	(7.1)		(100.0)				
記載なし																
(%)																
総計	303	429	1,160	4,435	14,554	45,989	93,912	78,173	28,854	7,308	2,673	277,790	32,318	310,108	10.71	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.2)	(16.6)	(33.8)	(28.1)	(10.4)	(2.6)	(1.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査