

表 3616 透析前ヘモグロビン濃度(g/dL) 原疾患別 (透析患者全体)

原疾患	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	84	87	292	1,201	4,135	13,944	29,659	25,184	9,260	2,264	840	86,950	10,020	96,970	10.75	1.25
(%)	(0.1)	(0.1)	(0.3)	(1.4)	(4.8)	(16.0)	(34.1)	(29.0)	(10.6)	(2.6)	(1.0)	(100.0)				
慢性腎盂腎炎	5	7	10	53	168	466	927	760	279	74	18	2,767	275	3,042	10.65	1.30
(%)	(0.2)	(0.3)	(0.4)	(1.9)	(6.1)	(16.8)	(33.5)	(27.5)	(10.1)	(2.7)	(0.7)	(100.0)				
急速進行性糸球体腎炎	1	6	9	45	122	376	730	600	241	54	21	2,205	289	2,494	10.68	1.30
(%)	(0.0)	(0.3)	(0.4)	(2.0)	(5.5)	(17.1)	(33.1)	(27.2)	(10.9)	(2.4)	(1.0)	(100.0)				
妊娠腎/妊娠中毒症	1	1	5	20	68	279	550	404	134	23	7	1,492	122	1,614	10.63	1.15
(%)	(0.1)	(0.1)	(0.3)	(1.3)	(4.6)	(18.7)	(36.9)	(27.1)	(9.0)	(1.5)	(0.5)	(100.0)				
その他の分類不能の腎炎	1	1	7	18	61	191	416	392	122	27	15	1,251	122	1,373	10.76	1.27
(%)	(0.1)	(0.1)	(0.6)	(1.4)	(4.9)	(15.3)	(33.3)	(31.3)	(9.8)	(2.2)	(1.2)	(100.0)				
多発性嚢胞腎	11	14	28	102	384	1,428	3,251	2,984	1,210	370	160	9,942	1,064	11,006	10.90	1.30
(%)	(0.1)	(0.1)	(0.3)	(1.0)	(3.9)	(14.4)	(32.7)	(30.0)	(12.2)	(3.7)	(1.6)	(100.0)				
腎硬化症	29	33	113	438	1,463	4,370	8,807	7,042	2,428	553	207	25,483	2,815	28,298	10.65	1.26
(%)	(0.1)	(0.1)	(0.4)	(1.7)	(5.7)	(17.1)	(34.6)	(27.6)	(9.5)	(2.2)	(0.8)	(100.0)				
悪性高血圧	6	3	4	40	115	358	765	674	227	63	31	2,286	287	2,573	10.75	1.34
(%)	(0.3)	(0.1)	(0.2)	(1.7)	(5.0)	(15.7)	(33.5)	(29.5)	(9.9)	(2.8)	(1.4)	(100.0)				
糖尿病性腎症	100	205	492	1,722	5,776	17,880	35,578	29,296	10,961	2,861	1,056	105,927	12,154	118,081	10.70	1.30
(%)	(0.1)	(0.2)	(0.5)	(1.6)	(5.5)	(16.9)	(33.6)	(27.7)	(10.3)	(2.7)	(1.0)	(100.0)				
SLE腎炎	3	3	9	42	123	321	675	530	209	55	23	1,993	243	2,236	10.67	1.33
(%)	(0.2)	(0.2)	(0.5)	(2.1)	(6.2)	(16.1)	(33.9)	(26.6)	(10.5)	(2.8)	(1.2)	(100.0)				
アミロイド腎			2	10	29	69	127	94	42	11	5	389	62	451	10.62	1.35
(%)			(0.5)	(2.6)	(7.5)	(17.7)	(32.6)	(24.2)	(10.8)	(2.8)	(1.3)	(100.0)				
痛風腎	1	1	2	16	34	174	329	303	111	24	11	1,006	107	1,113	10.79	1.23
(%)	(0.1)	(0.1)	(0.2)	(1.6)	(3.4)	(17.3)	(32.7)	(30.1)	(11.0)	(2.4)	(1.1)	(100.0)				
先天性代謝異常による腎不全		1	2	3	12	38	69	70	30	7	4	236	36	272	10.79	1.38
(%)		(0.4)	(0.8)	(1.3)	(5.1)	(16.1)	(29.2)	(29.7)	(12.7)	(3.0)	(1.7)	(100.0)				
腎・尿路結核			3	3	12	38	57	56	11	2	2	184	22	206	10.50	1.30
(%)			(1.6)	(1.6)	(6.5)	(20.7)	(31.0)	(30.4)	(6.0)	(1.1)	(1.1)	(100.0)				
腎・尿路結石	1		2	7	28	87	182	145	53	12	3	520	61	581	10.68	1.23
(%)	(0.2)		(0.4)	(1.3)	(5.4)	(16.7)	(35.0)	(27.9)	(10.2)	(2.3)	(0.6)	(100.0)				
腎・尿路腫瘍		3	7	20	57	142	280	196	64	15	4	788	98	886	10.49	1.30
(%)		(0.4)	(0.9)	(2.5)	(7.2)	(18.0)	(35.5)	(24.9)	(8.1)	(1.9)	(0.5)	(100.0)				
閉塞性尿路障害		1	2	9	36	102	205	193	63	27	1	639	84	723	10.75	1.27
(%)		(0.2)	(0.3)	(1.4)	(5.6)	(16.0)	(32.1)	(30.2)	(9.9)	(4.2)	(0.2)	(100.0)				
骨髄腫	1	2	3	20	19	54	55	51	18	8	2	233	43	276	10.20	1.66
(%)	(0.4)	(0.9)	(1.3)	(8.6)	(8.2)	(23.2)	(23.6)	(21.9)	(7.7)	(3.4)	(0.9)	(100.0)				
腎形成不全	1		1	9	27	80	191	174	72	24	9	588	61	649	10.88	1.30
(%)	(0.2)		(0.2)	(1.5)	(4.6)	(13.6)	(32.5)	(29.6)	(12.2)	(4.1)	(1.5)	(100.0)				
不明	42	48	115	462	1,426	4,151	8,206	6,629	2,394	611	176	24,260	3,284	27,544	10.64	1.30
(%)	(0.2)	(0.2)	(0.5)	(1.9)	(5.9)	(17.1)	(33.8)	(27.3)	(9.9)	(2.5)	(0.7)	(100.0)				
再導入	3	1	8	30	82	303	663	556	236	48	19	1,949	214	2,163	10.79	1.27
(%)	(0.2)	(0.1)	(0.4)	(1.5)	(4.2)	(15.5)	(34.0)	(28.5)	(12.1)	(2.5)	(1.0)	(100.0)				
その他	13	12	44	165	377	1,138	2,190	1,840	689	175	59	6,702	855	7,557	10.64	1.35
(%)	(0.2)	(0.2)	(0.7)	(2.5)	(5.6)	(17.0)	(32.7)	(27.5)	(10.3)	(2.6)	(0.9)	(100.0)				
合計	303	429	1,160	4,435	14,554	45,989	93,912	78,173	28,854	7,308	2,673	277,790	32,318	310,108	10.71	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.2)	(16.6)	(33.8)	(28.1)	(10.4)	(2.6)	(1.0)	(100.0)				
記載なし																
(%)																
総計	303	429	1,160	4,435	14,554	45,989	93,912	78,173	28,854	7,308	2,673	277,790	32,318	310,108	10.71	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.2)	(16.6)	(33.8)	(28.1)	(10.4)	(2.6)	(1.0)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査