

表 3618 透析前ヘモグロビン濃度(g/dL) 週透析回数(回/週)別 (透析患者全体)

| 週透析回数 | 5未満   | 5～    | 6～    | 7～    | 8～     | 9～     | 10～    | 11～    | 12～    | 13～    | 14～   | 合計      | 記載なし   | 総計      | 平均    | 標準偏差 |
|-------|-------|-------|-------|-------|--------|--------|--------|--------|--------|--------|-------|---------|--------|---------|-------|------|
| 1回    | 1     | 1     | 7     | 45    | 97     | 257    | 479    | 461    | 232    | 78     | 31    | 1,689   | 151    | 1,840   | 10.84 | 1.46 |
| (%)   | (0.1) | (0.1) | (0.4) | (2.7) | (5.7)  | (15.2) | (28.4) | (27.3) | (13.7) | (4.6)  | (1.8) | (100.0) |        |         |       |      |
| 2回    | 6     | 16    | 52    | 166   | 460    | 1,072  | 1,881  | 1,523  | 562    | 136    | 41    | 5,915   | 277    | 6,192   | 10.52 | 1.38 |
| (%)   | (0.1) | (0.3) | (0.9) | (2.8) | (7.8)  | (18.1) | (31.8) | (25.7) | (9.5)  | (2.3)  | (0.7) | (100.0) |        |         |       |      |
| 3回    | 287   | 395   | 1,059 | 4,044 | 13,465 | 43,209 | 88,730 | 73,508 | 26,925 | 6,762  | 2,467 | 260,851 | 9,788  | 270,639 | 10.71 | 1.28 |
| (%)   | (0.1) | (0.2) | (0.4) | (1.6) | (5.2)  | (16.6) | (34.0) | (28.2) | (10.3) | (2.6)  | (0.9) | (100.0) |        |         |       |      |
| 4回    |       | 1     | 1     | 16    | 50     | 135    | 249    | 290    | 142    | 42     | 24    | 950     | 44     | 994     | 10.99 | 1.39 |
| (%)   |       | (0.1) | (0.1) | (1.7) | (5.3)  | (14.2) | (26.2) | (30.5) | (14.9) | (4.4)  | (2.5) | (100.0) |        |         |       |      |
| 5回    | 1     |       |       | 1     | 2      | 7      | 23     | 26     | 12     | 4      | 3     | 79      | 10     | 89      | 11.16 | 1.66 |
| (%)   | (1.3) |       |       | (1.3) | (2.5)  | (8.9)  | (29.1) | (32.9) | (15.2) | (5.1)  | (3.8) | (100.0) |        |         |       |      |
| 6回    |       |       |       | 2     | 1      | 9      | 19     | 17     | 13     | 7      | 2     | 70      | 11     | 81      | 11.29 | 1.46 |
| (%)   |       |       |       | (2.9) | (1.4)  | (12.9) | (27.1) | (24.3) | (18.6) | (10.0) | (2.9) | (100.0) |        |         |       |      |
| 7回    |       |       |       |       | 2      | 4      | 2      | 5      | 7      | 5      | 2     | 27      | 3      | 30      | 11.93 | 1.84 |
| (%)   |       |       |       |       | (7.4)  | (14.8) | (7.4)  | (18.5) | (25.9) | (18.5) | (7.4) | (100.0) |        |         |       |      |
| 合計    | 295   | 413   | 1,119 | 4,274 | 14,077 | 44,693 | 91,383 | 75,830 | 27,893 | 7,034  | 2,570 | 269,581 | 10,284 | 279,865 | 10.71 | 1.28 |
| (%)   | (0.1) | (0.2) | (0.4) | (1.6) | (5.2)  | (16.6) | (33.9) | (28.1) | (10.3) | (2.6)  | (1.0) | (100.0) |        |         |       |      |
| 記載なし  | 8     | 16    | 41    | 161   | 477    | 1,296  | 2,529  | 2,343  | 961    | 274    | 103   | 8,209   | 22,034 | 30,243  | 10.75 | 1.37 |
| (%)   | (0.1) | (0.2) | (0.5) | (2.0) | (5.8)  | (15.8) | (30.8) | (28.5) | (11.7) | (3.3)  | (1.3) | (100.0) |        |         |       |      |
| 総計    | 303   | 429   | 1,160 | 4,435 | 14,554 | 45,989 | 93,912 | 78,173 | 28,854 | 7,308  | 2,673 | 277,790 | 32,318 | 310,108 | 10.71 | 1.28 |
| (%)   | (0.1) | (0.2) | (0.4) | (1.6) | (5.2)  | (16.6) | (33.8) | (28.1) | (10.4) | (2.6)  | (1.0) | (100.0) |        |         |       |      |
| 平均    | 2.98  | 2.96  | 2.94  | 2.95  | 2.96   | 2.97   | 2.97   | 2.97   | 2.97   | 2.97   | 2.98  | 2.97    | 2.95   | 2.97    |       |      |
| 標準偏差  | 0.22  | 0.22  | 0.26  | 0.29  | 0.26   | 0.23   | 0.22   | 0.23   | 0.26   | 0.30   | 0.31  | 0.23    | 0.33   | 0.24    |       |      |

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査