

表 3625 透析前ヘモグロビン濃度(g/dL) 性別（施設血液透析患者 透析回数週3回）

性別	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
男性	159	220	564	2,166	7,031	22,211	47,431	40,911	15,471	3,984	1,489	141,637	5,396	147,033	10.75	1.29
(%)	(0.1)	(0.2)	(0.4)	(1.5)	(5.0)	(15.7)	(33.5)	(28.9)	(10.9)	(2.8)	(1.1)	(100.0)				
女性	102	142	378	1,412	4,695	14,827	27,786	20,568	6,899	1,553	444	78,806	3,132	81,938	10.57	1.25
(%)	(0.1)	(0.2)	(0.5)	(1.8)	(6.0)	(18.8)	(35.3)	(26.1)	(8.8)	(2.0)	(0.6)	(100.0)				
合計	261	362	942	3,578	11,726	37,038	75,217	61,479	22,370	5,537	1,933	220,443	8,528	228,971	10.69	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.3)	(16.8)	(34.1)	(27.9)	(10.1)	(2.5)	(0.9)	(100.0)				
記載なし																
(%)																
総計	261	362	942	3,578	11,726	37,038	75,217	61,479	22,370	5,537	1,933	220,443	8,528	228,971	10.69	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.3)	(16.8)	(34.1)	(27.9)	(10.1)	(2.5)	(0.9)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査