

表 3628 透析前ヘモグロビン濃度(g/dL) 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	71	72	228	922	3,232	10,805	22,799	18,846	6,827	1,635	582	66,019	2,558	68,577	10.72	1.25
(%)	(0.1)	(0.1)	(0.3)	(1.4)	(4.9)	(16.4)	(34.5)	(28.5)	(10.3)	(2.5)	(0.9)	(100.0)				
慢性腎盂腎炎	4	3	10	44	122	383	698	572	204	48	12	2,100	65	2,165	10.61	1.29
(%)	(0.2)	(0.1)	(0.5)	(2.1)	(5.8)	(18.2)	(33.2)	(27.2)	(9.7)	(2.3)	(0.6)	(100.0)				
急速進行性糸球体腎炎	1	5	7	39	93	309	587	472	183	42	17	1,755	79	1,834	10.66	1.31
(%)	(0.1)	(0.3)	(0.4)	(2.2)	(5.3)	(17.6)	(33.4)	(26.9)	(10.4)	(2.4)	(1.0)	(100.0)				
妊娠腎/妊娠中毒症	1	1	3	15	50	184	389	276	91	16	4	1,030	34	1,064	10.62	1.16
(%)	(0.1)	(0.1)	(0.3)	(1.5)	(4.9)	(17.9)	(37.8)	(26.8)	(8.8)	(1.6)	(0.4)	(100.0)				
その他の分類不能の腎炎	1	1	7	14	45	153	296	296	78	19	8	918	31	949	10.70	1.28
(%)	(0.1)	(0.1)	(0.8)	(1.5)	(4.9)	(16.7)	(32.2)	(32.2)	(8.5)	(2.1)	(0.9)	(100.0)				
多発性嚢胞腎	11	12	21	87	316	1,119	2,642	2,352	945	282	122	7,909	289	8,198	10.88	1.30
(%)	(0.1)	(0.2)	(0.3)	(1.1)	(4.0)	(14.1)	(33.4)	(29.7)	(11.9)	(3.6)	(1.5)	(100.0)				
腎硬化症	27	30	92	363	1,181	3,574	7,262	5,670	1,952	422	154	20,727	733	21,460	10.63	1.26
(%)	(0.1)	(0.1)	(0.4)	(1.8)	(5.7)	(17.2)	(35.0)	(27.4)	(9.4)	(2.0)	(0.7)	(100.0)				
悪性高血圧	6	1	4	28	99	277	609	534	169	46	24	1,797	84	1,881	10.73	1.33
(%)	(0.3)	(0.1)	(0.2)	(1.6)	(5.5)	(15.4)	(33.9)	(29.7)	(9.4)	(2.6)	(1.3)	(100.0)				
糖尿病性腎症	84	175	417	1,417	4,793	14,878	29,524	24,063	8,863	2,253	785	87,252	3,255	90,507	10.68	1.29
(%)	(0.1)	(0.2)	(0.5)	(1.6)	(5.5)	(17.1)	(33.8)	(27.6)	(10.2)	(2.6)	(0.9)	(100.0)				
SLE腎炎	3	3	6	31	97	251	503	395	156	42	11	1,498	52	1,550	10.64	1.33
(%)	(0.2)	(0.2)	(0.4)	(2.1)	(6.5)	(16.8)	(33.6)	(26.4)	(10.4)	(2.8)	(0.7)	(100.0)				
アミロイド腎			1	9	19	54	96	75	35	9	5	303	22	325	10.68	1.37
(%)			(0.3)	(3.0)	(6.3)	(17.8)	(31.7)	(24.8)	(11.6)	(3.0)	(1.7)	(100.0)				
痛風腎	1	1	2	14	27	138	261	243	87	15	8	797	31	828	10.77	1.22
(%)	(0.1)	(0.1)	(0.3)	(1.8)	(3.4)	(17.3)	(32.7)	(30.5)	(10.9)	(1.9)	(1.0)	(100.0)				
先天性代謝異常による腎不全		1	1	3	10	30	51	49	18	4	2	169	8	177	10.66	1.32
(%)		(0.6)	(0.6)	(1.8)	(5.9)	(17.8)	(30.2)	(29.0)	(10.7)	(2.4)	(1.2)	(100.0)				
腎・尿路結核			3	2	9	33	50	45	9	2		153	5	158	10.46	1.23
(%)			(2.0)	(1.3)	(5.9)	(21.6)	(32.7)	(29.4)	(5.9)	(1.3)		(100.0)				
腎・尿路結石	1		2	6	23	68	147	108	43	10	3	411	19	430	10.66	1.27
(%)	(0.2)		(0.5)	(1.5)	(5.6)	(16.5)	(35.8)	(26.3)	(10.5)	(2.4)	(0.7)	(100.0)				
腎・尿路腫瘍		3	5	17	48	120	237	164	51	12	3	660	26	686	10.47	1.29
(%)		(0.5)	(0.8)	(2.6)	(7.3)	(18.2)	(35.9)	(24.8)	(7.7)	(1.8)	(0.5)	(100.0)				
閉塞性尿路障害		1	2	6	24	84	166	148	44	24		499	18	517	10.74	1.23
(%)		(0.2)	(0.4)	(1.2)	(4.8)	(16.8)	(33.3)	(29.7)	(8.8)	(4.8)		(100.0)				
骨髄腫	1	2	3	17	14	43	42	35	14	6	1	178	12	190	10.08	1.68
(%)	(0.6)	(1.1)	(1.7)	(9.6)	(7.9)	(24.2)	(23.6)	(19.7)	(7.9)	(3.4)	(0.6)	(100.0)				
腎形成不全	1			7	19	55	130	123	48	20	8	411	8	419	10.93	1.34
(%)	(0.2)			(1.7)	(4.6)	(13.4)	(31.6)	(29.9)	(11.7)	(4.9)	(1.9)	(100.0)				
不明	35	40	87	383	1,158	3,364	6,497	5,185	1,852	457	128	19,186	939	20,125	10.62	1.30
(%)	(0.2)	(0.2)	(0.5)	(2.0)	(6.0)	(17.5)	(33.9)	(27.0)	(9.7)	(2.4)	(0.7)	(100.0)				
再導入	3	1	5	22	52	214	484	385	165	33	15	1,379	38	1,417	10.78	1.27
(%)	(0.2)	(0.1)	(0.4)	(1.6)	(3.8)	(15.5)	(35.1)	(27.9)	(12.0)	(2.4)	(1.1)	(100.0)				
その他	10	10	36	132	295	902	1,747	1,443	536	140	41	5,292	222	5,514	10.62	1.34
(%)	(0.2)	(0.2)	(0.7)	(2.5)	(5.6)	(17.0)	(33.0)	(27.3)	(10.1)	(2.6)	(0.8)	(100.0)				
合計	261	362	942	3,578	11,726	37,038	75,217	61,479	22,370	5,537	1,933	220,443	8,528	228,971	10.69	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.3)	(16.8)	(34.1)	(27.9)	(10.1)	(2.5)	(0.9)	(100.0)				
記載なし																
(%)																
総計	261	362	942	3,578	11,726	37,038	75,217	61,479	22,370	5,537	1,933	220,443	8,528	228,971	10.69	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.3)	(16.8)	(34.1)	(27.9)	(10.1)	(2.5)	(0.9)	(100.0)				

数値下のかつこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査