

表 3630 透析前ヘモグロビン濃度(g/dL) 透析後体重(kg)別（施設血液透析患者 透析回数週3回）

透析後体重	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
20未満	1			1	6	12	32	23	1	5	1	82	2	84	10.53	1.47
(%)	(1.2)			(1.2)	(7.3)	(14.6)	(39.0)	(28.0)	(1.2)	(6.1)	(1.2)	(100.0)				
20～	1	2	5	36	66	152	249	174	71	18	6	780	22	802	10.43	1.47
(%)	(0.1)	(0.3)	(0.6)	(4.6)	(8.5)	(19.5)	(31.9)	(22.3)	(9.1)	(2.3)	(0.8)	(100.0)				
30～	26	36	112	460	1,447	3,808	6,102	4,547	1,673	409	133	18,753	364	19,117	10.49	1.33
(%)	(0.1)	(0.2)	(0.6)	(2.5)	(7.7)	(20.3)	(32.5)	(24.2)	(8.9)	(2.2)	(0.7)	(100.0)				
40～	82	111	284	1,136	3,678	10,944	20,242	15,307	5,368	1,264	362	58,778	1,031	59,809	10.58	1.28
(%)	(0.1)	(0.2)	(0.5)	(1.9)	(6.3)	(18.6)	(34.4)	(26.0)	(9.1)	(2.2)	(0.6)	(100.0)				
50～	81	124	276	1,095	3,571	11,897	24,639	20,050	7,112	1,637	499	70,981	1,128	72,109	10.69	1.25
(%)	(0.1)	(0.2)	(0.4)	(1.5)	(5.0)	(16.8)	(34.7)	(28.2)	(10.0)	(2.3)	(0.7)	(100.0)				
60～	45	54	158	514	1,815	6,551	14,920	12,912	4,766	1,204	474	43,413	669	44,082	10.80	1.25
(%)	(0.1)	(0.1)	(0.4)	(1.2)	(4.2)	(15.1)	(34.4)	(29.7)	(11.0)	(2.8)	(1.1)	(100.0)				
70～	15	17	58	190	688	2,191	5,646	5,216	2,076	590	249	16,936	248	17,184	10.90	1.28
(%)	(0.1)	(0.1)	(0.3)	(1.1)	(4.1)	(12.9)	(33.3)	(30.8)	(12.3)	(3.5)	(1.5)	(100.0)				
80～	3	5	20	61	200	789	1,943	1,868	765	231	104	5,989	92	6,081	10.97	1.28
(%)	(0.1)	(0.1)	(0.3)	(1.0)	(3.3)	(13.2)	(32.4)	(31.2)	(12.8)	(3.9)	(1.7)	(100.0)				
90～	2	3	4	22	82	273	683	683	269	87	59	2,167	28	2,195	10.99	1.33
(%)	(0.1)	(0.1)	(0.2)	(1.0)	(3.8)	(12.6)	(31.5)	(31.5)	(12.4)	(4.0)	(2.7)	(100.0)				
100～	1	1	4	8	27	93	247	240	82	33	13	749	6	755	10.93	1.31
(%)	(0.1)	(0.1)	(0.5)	(1.1)	(3.6)	(12.4)	(33.0)	(32.0)	(10.9)	(4.4)	(1.7)	(100.0)				
110～				5	16	43	95	88	31	6	12	296	3	299	10.90	1.40
(%)				(1.7)	(5.4)	(14.5)	(32.1)	(29.7)	(10.5)	(2.0)	(4.1)	(100.0)				
120～			1	4	9	33	57	46	22	10	5	187	2	189	10.85	1.45
(%)			(0.5)	(2.1)	(4.8)	(17.6)	(30.5)	(24.6)	(11.8)	(5.3)	(2.7)	(100.0)				
合計	257	353	922	3,532	11,605	36,786	74,855	61,154	22,236	5,494	1,917	219,111	3,595	222,706	10.69	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.3)	(16.8)	(34.2)	(27.9)	(10.1)	(2.5)	(0.9)	(100.0)				
記載なし	4	9	20	46	121	252	362	325	134	43	16	1,332	4,933	6,265	10.45	1.57
(%)	(0.3)	(0.7)	(1.5)	(3.5)	(9.1)	(18.9)	(27.2)	(24.4)	(10.1)	(3.2)	(1.2)	(100.0)				
総計	261	362	942	3,578	11,726	37,038	75,217	61,479	22,370	5,537	1,933	220,443	8,528	228,971	10.69	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.3)	(16.8)	(34.1)	(27.9)	(10.1)	(2.5)	(0.9)	(100.0)				
平均	53.16	52.94	53.17	52.38	52.86	53.91	55.42	56.35	56.76	57.75	60.26	55.46	54.30	55.45		
標準偏差	12.11	11.89	12.66	13.14	12.56	12.49	12.65	12.97	13.21	14.14	15.40	12.90	12.57	12.90		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査