

表 3641 透析前ヘモグロビン濃度(g/dL) 医療機関別（血液透析濾過患者 透析回数週3回）

医療機関	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
国公立大学		1		2	5	6	12	9	5	1	1	42		42	10.51	1.71
(%)		(2.4)		(4.8)	(11.9)	(14.3)	(28.6)	(21.4)	(11.9)	(2.4)	(2.4)	(100.0)				
私立大学				3	11	26	55	50	20	3	3	171	2	173	10.77	1.25
(%)				(1.8)	(6.4)	(15.2)	(32.2)	(29.2)	(11.7)	(1.8)	(1.8)	(100.0)				
国立						7	7	4	4			22	14	36	10.69	1.08
(%)						(31.8)	(31.8)	(18.2)	(18.2)			(100.0)				
県市町村立		1	9	32	95	267	515	501	204	54	20	1,698	75	1,773	10.78	1.33
(%)		(0.1)	(0.5)	(1.9)	(5.6)	(15.7)	(30.3)	(29.5)	(12.0)	(3.2)	(1.2)	(100.0)				
社会保険		1	2	5	13	50	98	67	27	7	3	273	46	319	10.65	1.33
(%)		(0.4)	(0.7)	(1.8)	(4.8)	(18.3)	(35.9)	(24.5)	(9.9)	(2.6)	(1.1)	(100.0)				
厚生連			5	19	42	151	301	286	127	29	15	975	38	1,013	10.85	1.34
(%)			(0.5)	(1.9)	(4.3)	(15.5)	(30.9)	(29.3)	(13.0)	(3.0)	(1.5)	(100.0)				
その他公的		1	1	13	39	90	161	227	79	25	11	647	47	694	10.93	1.41
(%)		(0.2)	(0.2)	(2.0)	(6.0)	(13.9)	(24.9)	(35.1)	(12.2)	(3.9)	(1.7)	(100.0)				
私立総合	1	1	5	17	44	169	335	335	124	40	9	1,080	13	1,093	10.82	1.26
(%)	(0.1)	(0.1)	(0.5)	(1.6)	(4.1)	(15.6)	(31.0)	(31.0)	(11.5)	(3.7)	(0.8)	(100.0)				
私立	5	6	33	130	442	1,523	3,186	2,739	1,143	303	132	9,642	373	10,015	10.82	1.29
(%)	(0.1)	(0.1)	(0.3)	(1.3)	(4.6)	(15.8)	(33.0)	(28.4)	(11.9)	(3.1)	(1.4)	(100.0)				
私立診療所	20	20	59	224	946	3,561	8,277	7,359	2,649	723	329	24,167	621	24,788	10.85	1.24
(%)	(0.1)	(0.1)	(0.2)	(0.9)	(3.9)	(14.7)	(34.2)	(30.5)	(11.0)	(3.0)	(1.4)	(100.0)				
総計	26	31	114	445	1,637	5,850	12,947	11,577	4,382	1,185	523	38,717	1,229	39,946	10.84	1.26
(%)	(0.1)	(0.1)	(0.3)	(1.1)	(4.2)	(15.1)	(33.4)	(29.9)	(11.3)	(3.1)	(1.4)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査