

表 3644 透析前ヘモグロビン濃度(g/dL) HDL-C濃度(mg/dL)別 (血液透析濾過患者 透析回数週3回)

HDL-C濃度	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
20未満	1	2	2	7	16	11	17	14	6	4	1	81		81	9.88	2.02
(%)	(1.2)	(2.5)	(2.5)	(8.6)	(19.8)	(13.6)	(21.0)	(17.3)	(7.4)	(4.9)	(1.2)	(100.0)				
20～	2	1	10	35	115	306	577	510	212	61	28	1,857	17	1,874	10.74	1.37
(%)	(0.1)	(0.1)	(0.5)	(1.9)	(6.2)	(16.5)	(31.1)	(27.5)	(11.4)	(3.3)	(1.5)	(100.0)				
30～	1	4	22	89	289	952	2,000	1,754	659	211	101	6,082	49	6,131	10.82	1.31
(%)	(0.0)	(0.1)	(0.4)	(1.5)	(4.8)	(15.7)	(32.9)	(28.8)	(10.8)	(3.5)	(1.7)	(100.0)				
40～	3	10	23	89	294	1,041	2,480	2,200	821	248	101	7,310	50	7,360	10.87	1.27
(%)	(0.0)	(0.1)	(0.3)	(1.2)	(4.0)	(14.2)	(33.9)	(30.1)	(11.2)	(3.4)	(1.4)	(100.0)				
50～	1	3	12	45	211	887	1,938	1,895	679	179	78	5,928	43	5,971	10.89	1.22
(%)	(0.0)	(0.1)	(0.2)	(0.8)	(3.6)	(15.0)	(32.7)	(32.0)	(11.5)	(3.0)	(1.3)	(100.0)				
60～	5	3	6	26	114	518	1,347	1,223	455	101	35	3,833	24	3,857	10.89	1.15
(%)	(0.1)	(0.1)	(0.2)	(0.7)	(3.0)	(13.5)	(35.1)	(31.9)	(11.9)	(2.6)	(0.9)	(100.0)				
70～	2	1	2	11	71	249	719	618	269	52	15	2,009	14	2,023	10.90	1.14
(%)	(0.1)	(0.0)	(0.1)	(0.5)	(3.5)	(12.4)	(35.8)	(30.8)	(13.4)	(2.6)	(0.7)	(100.0)				
80～	2		3	3	26	136	378	312	115	20	10	1,005	8	1,013	10.90	1.17
(%)	(0.2)		(0.3)	(0.3)	(2.6)	(13.5)	(37.6)	(31.0)	(11.4)	(2.0)	(1.0)	(100.0)				
90～				3	12	68	140	148	41	11	5	428	3	431	10.89	1.12
(%)				(0.7)	(2.8)	(15.9)	(32.7)	(34.6)	(9.6)	(2.6)	(1.2)	(100.0)				
100～		1	2	2	9	50	143	123	37	2	5	374	9	383	10.83	1.12
(%)		(0.3)	(0.5)	(0.5)	(2.4)	(13.4)	(38.2)	(32.9)	(9.9)	(0.5)	(1.3)	(100.0)				
合計	17	25	82	310	1,157	4,218	9,739	8,797	3,294	889	379	28,907	217	29,124	10.86	1.25
(%)	(0.1)	(0.1)	(0.3)	(1.1)	(4.0)	(14.6)	(33.7)	(30.4)	(11.4)	(3.1)	(1.3)	(100.0)				
記載なし	9	6	32	135	480	1,632	3,208	2,780	1,088	296	144	9,810	1,012	10,822	10.78	1.30
(%)	(0.1)	(0.1)	(0.3)	(1.4)	(4.9)	(16.6)	(32.7)	(28.3)	(11.1)	(3.0)	(1.5)	(100.0)				
総計	26	31	114	445	1,637	5,850	12,947	11,577	4,382	1,185	523	38,717	1,229	39,946	10.84	1.26
(%)	(0.1)	(0.1)	(0.3)	(1.1)	(4.2)	(15.1)	(33.4)	(29.9)	(11.3)	(3.1)	(1.4)	(100.0)				
平均	53.65	46.36	45.56	44.35	47.15	50.06	51.46	51.54	51.23	48.25	48.12	50.84	52.49	50.85		
標準偏差	20.22	19.45	18.38	15.92	16.35	17.25	17.43	17.09	16.75	14.98	16.69	17.14	22.02	17.18		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査