

表 3653 ヘモグロビン濃度(g/dL) 医療機関別（腹膜透析患者）

医療機関	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
国公立大学	2	2		7	27	56	156	156	73	13	4	496	217	713	10.85	1.30
(%)	(0.4)	(0.4)		(1.4)	(5.4)	(11.3)	(31.5)	(31.5)	(14.7)	(2.6)	(0.8)	(100.0)				
私立大学	3		2	9	35	110	198	204	79	27	2	669	464	1,133	10.75	1.37
(%)	(0.4)		(0.3)	(1.3)	(5.2)	(16.4)	(29.6)	(30.5)	(11.8)	(4.0)	(0.3)	(100.0)				
国立	2	3	3	11	13	22	35	29	13	4	2	137	65	202	10.09	1.92
(%)	(1.5)	(2.2)	(2.2)	(8.0)	(9.5)	(16.1)	(25.5)	(21.2)	(9.5)	(2.9)	(1.5)	(100.0)				
県市町村立			5	25	66	150	311	294	124	40	14	1,029	557	1,586	10.78	1.37
(%)			(0.5)	(2.4)	(6.4)	(14.6)	(30.2)	(28.6)	(12.1)	(3.9)	(1.4)	(100.0)				
社会保険			1	2	15	42	64	52	19	5	4	204	283	487	10.64	1.35
(%)			(0.5)	(1.0)	(7.4)	(20.6)	(31.4)	(25.5)	(9.3)	(2.5)	(2.0)	(100.0)				
厚生連		2	4	13	31	91	121	121	69	19	9	480	146	626	10.76	1.52
(%)		(0.4)	(0.8)	(2.7)	(6.5)	(19.0)	(25.2)	(25.2)	(14.4)	(4.0)	(1.9)	(100.0)				
その他公的		1	3	17	48	147	259	283	125	46	19	948	371	1,319	10.92	1.43
(%)		(0.1)	(0.3)	(1.8)	(5.1)	(15.5)	(27.3)	(29.9)	(13.2)	(4.9)	(2.0)	(100.0)				
私立総合		4	2	4	10	33	58	56	36	6		209	168	377	10.70	1.50
(%)		(1.9)	(1.0)	(1.9)	(4.8)	(15.8)	(27.8)	(26.8)	(17.2)	(2.9)		(100.0)				
私立		1	6	31	69	163	300	305	161	52	21	1,109	580	1,689	10.87	1.48
(%)		(0.1)	(0.5)	(2.8)	(6.2)	(14.7)	(27.1)	(27.5)	(14.5)	(4.7)	(1.9)	(100.0)				
私立診療所		1	2	5	30	89	204	209	97	25	4	666	143	809	10.93	1.23
(%)		(0.2)	(0.3)	(0.8)	(4.5)	(13.4)	(30.6)	(31.4)	(14.6)	(3.8)	(0.6)	(100.0)				
総計	7	14	28	124	344	903	1,706	1,709	796	237	79	5,947	2,994	8,941	10.81	1.42
(%)	(0.1)	(0.2)	(0.5)	(2.1)	(5.8)	(15.2)	(28.7)	(28.7)	(13.4)	(4.0)	(1.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査