

表 3663 総コレステロール濃度(mg/dL) 併用の状況別（透析患者全体）

併用の状況	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	7,774	22,537	41,701	47,798	39,333	24,663	13,142	5,744	2,239	1,432	206,363	94,431	300,794	155.37	35.46
(%)	(3.8)	(10.9)	(20.2)	(23.2)	(19.1)	(12.0)	(6.4)	(2.8)	(1.1)	(0.7)	(100.0)				
PD単独	33	101	280	557	634	678	463	271	136	140	3,293	3,895	7,188	183.26	41.29
(%)	(1.0)	(3.1)	(8.5)	(16.9)	(19.3)	(20.6)	(14.1)	(8.2)	(4.1)	(4.3)	(100.0)				
HD(F)単独.PDカテあり	4	18	35	42	27	15	15	5	2	3	166	103	269	158.14	37.19
(%)	(2.4)	(10.8)	(21.1)	(25.3)	(16.3)	(9.0)	(9.0)	(3.0)	(1.2)	(1.8)	(100.0)				
PD.HD併用週1 HD(F)	12	56	155	216	196	135	88	40	14	9	921	623	1,544	164.96	34.71
(%)	(1.3)	(6.1)	(16.8)	(23.5)	(21.3)	(14.7)	(9.6)	(4.3)	(1.5)	(1.0)	(100.0)				
PD.HD併用週2 HD(F)	2	4	12	28	28	19	12	9	8	1	123	54	177	174.49	37.50
(%)	(1.6)	(3.3)	(9.8)	(22.8)	(22.8)	(15.4)	(9.8)	(7.3)	(6.5)	(0.8)	(100.0)				
PD.HD併用週3 HD(F)	1	4	5	3	9	5	1	1			29	11	40	157.76	33.01
(%)	(3.4)	(13.8)	(17.2)	(10.3)	(31.0)	(17.2)	(3.4)	(3.4)			(100.0)				
PD.HD併用週4 HD(F)			1								1		1	131.00	
(%)			(100.0)								(100.0)				
PD.HD併用その他	3	1	6	11	14	10	2	1	1		49	24	73	162.88	32.95
(%)	(6.1)	(2.0)	(12.2)	(22.4)	(28.6)	(20.4)	(4.1)	(2.0)	(2.0)		(100.0)				
合計	7,829	22,721	42,195	48,655	40,241	25,525	13,723	6,071	2,400	1,585	210,945	99,141	310,086	155.86	35.74
(%)	(3.7)	(10.8)	(20.0)	(23.1)	(19.1)	(12.1)	(6.5)	(2.9)	(1.1)	(0.8)	(100.0)				
不明	1	1	6	4	2					1	15	7	22	148.53	42.92
(%)	(6.7)	(6.7)	(40.0)	(26.7)	(13.3)					(6.7)	(100.0)				
記載なし (%)															
総計	7,830	22,722	42,201	48,659	40,243	25,525	13,723	6,071	2,400	1,586	210,960	99,148	310,108	155.86	35.74
(%)	(3.7)	(10.8)	(20.0)	(23.1)	(19.1)	(12.1)	(6.5)	(2.9)	(1.1)	(0.8)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査