

表 3665 総コレステロール濃度(mg/dL) 年齢(歳)別 (透析患者全体)

年齢	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	合計	記載なし	総計	平均	標準偏差
15歳未満	1	2	7	3	7	7	7	7	4	8	53	64	117	204.85	68.75
(%)	(1.9)	(3.8)	(13.2)	(5.7)	(13.2)	(13.2)	(13.2)	(13.2)	(7.5)	(15.1)	(100.0)				
15歳～	30	91	151	144	129	73	44	11	5	5	683	354	1,037	151.74	34.95
(%)	(4.4)	(13.3)	(22.1)	(21.1)	(18.9)	(10.7)	(6.4)	(1.6)	(0.7)	(0.7)	(100.0)				
30歳～	220	838	1,759	2,264	2,062	1,366	731	325	123	108	9,796	4,589	14,385	160.72	35.82
(%)	(2.2)	(8.6)	(18.0)	(23.1)	(21.0)	(13.9)	(7.5)	(3.3)	(1.3)	(1.1)	(100.0)				
45歳～	1,096	3,346	6,688	8,601	7,816	5,378	3,238	1,464	622	391	38,640	17,928	56,568	161.46	36.72
(%)	(2.8)	(8.7)	(17.3)	(22.3)	(20.2)	(13.9)	(8.4)	(3.8)	(1.6)	(1.0)	(100.0)				
60歳～	3,630	10,335	19,179	22,190	18,559	11,902	6,460	2,831	1,092	744	96,922	45,056	141,978	156.15	35.86
(%)	(3.7)	(10.7)	(19.8)	(22.9)	(19.1)	(12.3)	(6.7)	(2.9)	(1.1)	(0.8)	(100.0)				
75歳～	2,697	7,680	13,687	14,655	11,043	6,450	3,090	1,371	519	313	61,505	29,492	90,997	151.37	34.26
(%)	(4.4)	(12.5)	(22.3)	(23.8)	(18.0)	(10.5)	(5.0)	(2.2)	(0.8)	(0.5)	(100.0)				
90歳～	156	428	729	801	624	348	153	62	35	17	3,353	1,652	5,005	151.00	34.16
(%)	(4.7)	(12.8)	(21.7)	(23.9)	(18.6)	(10.4)	(4.6)	(1.8)	(1.0)	(0.5)	(100.0)				
合計	7,830	22,720	42,200	48,658	40,240	25,524	13,723	6,071	2,400	1,586	210,952	99,135	310,087	155.86	35.74
(%)	(3.7)	(10.8)	(20.0)	(23.1)	(19.1)	(12.1)	(6.5)	(2.9)	(1.1)	(0.8)	(100.0)				
不明		2	1	1	3	1					8	13	21	150.13	25.84
(%)		(25.0)	(12.5)	(12.5)	(37.5)	(12.5)					(100.0)				
記載なし															
(%)															
総計	7,830	22,722	42,201	48,659	40,243	25,525	13,723	6,071	2,400	1,586	210,960	99,148	310,108	155.86	35.74
(%)	(3.7)	(10.8)	(20.0)	(23.1)	(19.1)	(12.1)	(6.5)	(2.9)	(1.1)	(0.8)	(100.0)				
平均	69.62	69.09	68.52	67.78	66.92	66.19	65.35	65.23	65.06	63.97	67.49	67.64	67.54		
標準偏差	11.84	12.13	12.33	12.43	12.51	12.51	12.35	12.37	12.57	13.18	12.44	12.60	12.49		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査