

表 3667 総コレステロール濃度(mg/dL) 原疾患別 (透析患者全体)

原疾患	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	1,813	5,740	12,058	15,248	13,539	9,002	5,069	2,189	876	522	66,056	30,914	96,970	159.83	35.38
(%)	(2.7)	(8.7)	(18.3)	(23.1)	(20.5)	(13.6)	(7.7)	(3.3)	(1.3)	(0.8)	(100.0)				
慢性腎盂腎炎	56	161	372	444	415	301	172	99	30	21	2,071	971	3,042	162.74	36.83
(%)	(2.7)	(7.8)	(18.0)	(21.4)	(20.0)	(14.5)	(8.3)	(4.8)	(1.4)	(1.0)	(100.0)				
急速進行性糸球体腎炎	32	96	237	338	344	277	156	101	49	48	1,678	816	2,494	170.89	40.91
(%)	(1.9)	(5.7)	(14.1)	(20.1)	(20.5)	(16.5)	(9.3)	(6.0)	(2.9)	(2.9)	(100.0)				
妊娠腎/妊娠中毒症	10	47	137	220	250	224	133	74	26	15	1,136	478	1,614	173.31	35.19
(%)	(0.9)	(4.1)	(12.1)	(19.4)	(22.0)	(19.7)	(11.7)	(6.5)	(2.3)	(1.3)	(100.0)				
その他の分類不能の腎炎	29	82	169	196	199	133	89	29	8	11	945	428	1,373	161.28	37.85
(%)	(3.1)	(8.7)	(17.9)	(20.7)	(21.1)	(14.1)	(9.4)	(3.1)	(0.8)	(1.2)	(100.0)				
多発性嚢胞腎	187	638	1,404	1,810	1,654	1,068	530	256	76	52	7,675	3,331	11,006	159.49	34.11
(%)	(2.4)	(8.3)	(18.3)	(23.6)	(21.6)	(13.9)	(6.9)	(3.3)	(1.0)	(0.7)	(100.0)				
腎硬化症	576	1,970	4,128	4,743	3,873	2,397	1,184	506	200	129	19,706	8,592	28,298	155.77	34.09
(%)	(2.9)	(10.0)	(20.9)	(24.1)	(19.7)	(12.2)	(6.0)	(2.6)	(1.0)	(0.7)	(100.0)				
悪性高血圧	40	154	315	385	371	201	135	62	27	16	1,706	867	2,573	160.60	36.86
(%)	(2.3)	(9.0)	(18.5)	(22.6)	(21.7)	(11.8)	(7.9)	(3.6)	(1.6)	(0.9)	(100.0)				
糖尿病性腎症	4,124	11,007	17,904	18,584	13,853	8,028	4,078	1,788	706	487	80,559	37,522	118,081	150.29	35.25
(%)	(5.1)	(13.7)	(22.2)	(23.1)	(17.2)	(10.0)	(5.1)	(2.2)	(0.9)	(0.6)	(100.0)				
SLE腎炎	33	100	228	296	314	222	172	104	46	27	1,542	694	2,236	169.97	40.55
(%)	(2.1)	(6.5)	(14.8)	(19.2)	(20.4)	(14.4)	(11.2)	(6.7)	(3.0)	(1.8)	(100.0)				
アミロイド腎	19	28	61	57	53	36	24	17	5	4	304	147	451	158.66	41.27
(%)	(6.3)	(9.2)	(20.1)	(18.8)	(17.4)	(11.8)	(7.9)	(5.6)	(1.6)	(1.3)	(100.0)				
痛風腎	24	66	175	186	138	109	47	21	8	5	779	334	1,113	156.29	34.69
(%)	(3.1)	(8.5)	(22.5)	(23.9)	(17.7)	(14.0)	(6.0)	(2.7)	(1.0)	(0.6)	(100.0)				
先天性代謝異常による腎不全	8	18	32	33	30	24	18	9	2	4	178	94	272	162.44	42.57
(%)	(4.5)	(10.1)	(18.0)	(18.5)	(16.9)	(13.5)	(10.1)	(5.1)	(1.1)	(2.2)	(100.0)				
腎・尿路結核	3	13	19	39	24	14	12	4	1	2	131	75	206	159.78	35.13
(%)	(2.3)	(9.9)	(14.5)	(29.8)	(18.3)	(10.7)	(9.2)	(3.1)	(0.8)	(1.5)	(100.0)				
腎・尿路結石	12	39	72	105	75	54	24	9	5	1	396	185	581	156.10	33.29
(%)	(3.0)	(9.8)	(18.2)	(26.5)	(18.9)	(13.6)	(6.1)	(2.3)	(1.3)	(0.3)	(100.0)				
腎・尿路腫瘍	28	65	142	152	108	66	35	17	7	3	623	263	886	153.02	35.17
(%)	(4.5)	(10.4)	(22.8)	(24.4)	(17.3)	(10.6)	(5.6)	(2.7)	(1.1)	(0.5)	(100.0)				
閉塞性尿路障害	11	40	88	81	106	74	42	25	6	5	478	245	723	163.86	37.54
(%)	(2.3)	(8.4)	(18.4)	(16.9)	(22.2)	(15.5)	(8.8)	(5.2)	(1.3)	(1.0)	(100.0)				
骨髄腫	5	24	19	42	29	32	12	10	2	6	181	95	276	164.50	42.25
(%)	(2.8)	(13.3)	(10.5)	(23.2)	(16.0)	(17.7)	(6.6)	(5.5)	(1.1)	(3.3)	(100.0)				
腎形成不全	9	31	84	86	83	69	36	22	9	3	432	217	649	163.34	35.34
(%)	(2.1)	(7.2)	(19.4)	(19.9)	(19.2)	(16.0)	(8.3)	(5.1)	(2.1)	(0.7)	(100.0)				
不明	583	1,804	3,381	4,164	3,531	2,290	1,249	520	205	142	17,869	9,675	27,544	157.56	35.60
(%)	(3.3)	(10.1)	(18.9)	(23.3)	(19.8)	(12.8)	(7.0)	(2.9)	(1.1)	(0.8)	(100.0)				
再導入	38	123	284	356	292	220	125	51	27	17	1,533	630	2,163	161.65	36.94
(%)	(2.5)	(8.0)	(18.5)	(23.2)	(19.0)	(14.4)	(8.2)	(3.3)	(1.8)	(1.1)	(100.0)				
その他	190	476	892	1,094	962	684	381	158	79	66	4,982	2,575	7,557	159.35	37.76
(%)	(3.8)	(9.6)	(17.9)	(22.0)	(19.3)	(13.7)	(7.6)	(3.2)	(1.6)	(1.3)	(100.0)				
合計	7,830	22,722	42,201	48,659	40,243	25,525	13,723	6,071	2,400	1,586	210,960	99,148	310,108	155.86	35.74
(%)	(3.7)	(10.8)	(20.0)	(23.1)	(19.1)	(12.1)	(6.5)	(2.9)	(1.1)	(0.8)	(100.0)				
記載なし															
(%)															
総計	7,830	22,722	42,201	48,659	40,243	25,525	13,723	6,071	2,400	1,586	210,960	99,148	310,108	155.86	35.74
(%)	(3.7)	(10.8)	(20.0)	(23.1)	(19.1)	(12.1)	(6.5)	(2.9)	(1.1)	(0.8)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査