

表 3678 総コレステロール濃度(mg/dL) 原疾患別 (施設血液透析患者 透析回数週3回)

原疾患	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	1,446	4,499	9,355	11,682	10,221	6,612	3,594	1,501	599	350	49,859	18,718	68,577	158.57	34.91
(%)	(2.9)	(9.0)	(18.8)	(23.4)	(20.5)	(13.3)	(7.2)	(3.0)	(1.2)	(0.7)	(100.0)				
慢性腎盂腎炎	42	123	283	343	285	226	125	67	21	12	1,527	638	2,165	161.53	36.22
(%)	(2.8)	(8.1)	(18.5)	(22.5)	(18.7)	(14.8)	(8.2)	(4.4)	(1.4)	(0.8)	(100.0)				
急速進行性糸球体腎炎	25	78	198	255	285	218	119	81	33	38	1,330	504	1,834	170.12	40.46
(%)	(1.9)	(5.9)	(14.9)	(19.2)	(21.4)	(16.4)	(8.9)	(6.1)	(2.5)	(2.9)	(100.0)				
妊娠腎/妊娠中毒症	6	35	95	152	168	156	78	47	16	10	763	301	1,064	172.09	34.73
(%)	(0.8)	(4.6)	(12.5)	(19.9)	(22.0)	(20.4)	(10.2)	(6.2)	(2.1)	(1.3)	(100.0)				
その他の分類不能の腎炎	20	62	135	156	149	85	62	20	5	9	703	246	949	159.78	37.85
(%)	(2.8)	(8.8)	(19.2)	(22.2)	(21.2)	(12.1)	(8.8)	(2.8)	(0.7)	(1.3)	(100.0)				
多発性嚢胞腎	147	516	1,126	1,423	1,313	849	407	192	62	42	6,077	2,121	8,198	159.18	34.09
(%)	(2.4)	(8.5)	(18.5)	(23.4)	(21.6)	(14.0)	(6.7)	(3.2)	(1.0)	(0.7)	(100.0)				
腎硬化症	478	1,664	3,462	3,921	3,147	1,898	902	377	153	86	16,088	5,372	21,460	154.68	33.47
(%)	(3.0)	(10.3)	(21.5)	(24.4)	(19.6)	(11.8)	(5.6)	(2.3)	(1.0)	(0.5)	(100.0)				
悪性高血圧	32	127	253	298	289	151	103	44	23	10	1,330	551	1,881	159.45	36.39
(%)	(2.4)	(9.5)	(19.0)	(22.4)	(21.7)	(11.4)	(7.7)	(3.3)	(1.7)	(0.8)	(100.0)				
糖尿病性腎症	3,457	9,241	15,012	15,294	11,277	6,487	3,204	1,447	553	381	66,353	24,154	90,507	149.68	35.03
(%)	(5.2)	(13.9)	(22.6)	(23.0)	(17.0)	(9.8)	(4.8)	(2.2)	(0.8)	(0.6)	(100.0)				
SLE腎炎	26	74	172	235	236	160	124	80	30	12	1,149	401	1,550	168.39	38.96
(%)	(2.3)	(6.4)	(15.0)	(20.5)	(20.5)	(13.9)	(10.8)	(7.0)	(2.6)	(1.0)	(100.0)				
アミロイド腎	16	25	50	44	43	30	17	10	2	3	240	85	325	155.99	40.31
(%)	(6.7)	(10.4)	(20.8)	(18.3)	(17.9)	(12.5)	(7.1)	(4.2)	(0.8)	(1.3)	(100.0)				
痛風腎	18	51	145	140	105	92	38	13	4	3	609	219	828	155.53	33.97
(%)	(3.0)	(8.4)	(23.8)	(23.0)	(17.2)	(15.1)	(6.2)	(2.1)	(0.7)	(0.5)	(100.0)				
先天性代謝異常による腎不全	8	13	24	21	19	19	12	5	2	1	124	53	177	158.15	39.77
(%)	(6.5)	(10.5)	(19.4)	(16.9)	(15.3)	(15.3)	(9.7)	(4.0)	(1.6)	(0.8)	(100.0)				
腎・尿路結核	3	11	18	31	18	11	9	3	1	1	106	52	158	157.35	34.60
(%)	(2.8)	(10.4)	(17.0)	(29.2)	(17.0)	(10.4)	(8.5)	(2.8)	(0.9)	(0.9)	(100.0)				
腎・尿路結石	11	32	60	82	62	44	17	5	5		318	112	430	154.70	32.70
(%)	(3.5)	(10.1)	(18.9)	(25.8)	(19.5)	(13.8)	(5.3)	(1.6)	(1.6)		(100.0)				
腎・尿路腫瘍	25	57	120	129	85	60	29	11	7	3	526	160	686	152.56	35.35
(%)	(4.8)	(10.8)	(22.8)	(24.5)	(16.2)	(11.4)	(5.5)	(2.1)	(1.3)	(0.6)	(100.0)				
閉塞性尿路障害	9	33	74	70	82	57	34	17	4	4	384	133	517	162.36	36.52
(%)	(2.3)	(8.6)	(19.3)	(18.2)	(21.4)	(14.8)	(8.9)	(4.4)	(1.0)	(1.0)	(100.0)				
骨髄腫	4	20	13	31	21	29	8	5	1	3	135	55	190	161.72	39.51
(%)	(3.0)	(14.8)	(9.6)	(23.0)	(15.6)	(21.5)	(5.9)	(3.7)	(0.7)	(2.2)	(100.0)				
腎形成不全	6	21	56	67	55	43	27	15	6	1	297	122	419	162.84	34.60
(%)	(2.0)	(7.1)	(18.9)	(22.6)	(18.5)	(14.5)	(9.1)	(5.1)	(2.0)	(0.3)	(100.0)				
不明	489	1,462	2,725	3,341	2,765	1,713	945	379	132	90	14,041	6,084	20,125	156.15	34.94
(%)	(3.5)	(10.4)	(19.4)	(23.8)	(19.7)	(12.2)	(6.7)	(2.7)	(0.9)	(0.6)	(100.0)				
再導入	29	91	198	244	208	166	79	36	17	8	1,076	341	1,417	160.91	35.63
(%)	(2.7)	(8.5)	(18.4)	(22.7)	(19.3)	(15.4)	(7.3)	(3.3)	(1.6)	(0.7)	(100.0)				
その他	157	385	741	867	750	513	276	120	57	44	3,910	1,604	5,514	157.80	37.11
(%)	(4.0)	(9.8)	(19.0)	(22.2)	(19.2)	(13.1)	(7.1)	(3.1)	(1.5)	(1.1)	(100.0)				
合計	6,454	18,620	34,315	38,826	31,583	19,619	10,209	4,475	1,733	1,111	166,945	62,026	228,971	154.71	35.28
(%)	(3.9)	(11.2)	(20.6)	(23.3)	(18.9)	(11.8)	(6.1)	(2.7)	(1.0)	(0.7)	(100.0)				
記載なし															
(%)															
総計	6,454	18,620	34,315	38,826	31,583	19,619	10,209	4,475	1,733	1,111	166,945	62,026	228,971	154.71	35.28
(%)	(3.9)	(11.2)	(20.6)	(23.3)	(18.9)	(11.8)	(6.1)	(2.7)	(1.0)	(0.7)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査