

表 3689 総コレステロール濃度(mg/dL) 原疾患別(血液透析濾過患者 透析回数週3回)

原疾患	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	270	905	1,975	2,582	2,370	1,650	979	456	161	89	11,437	3,639	15,076	162.02	35.33
(%)	(2.4)	(7.9)	(17.3)	(22.6)	(20.7)	(14.4)	(8.6)	(4.0)	(1.4)	(0.8)	(100.0)				
慢性腎盂腎炎	4	24	62	69	89	47	24	18	5	2	344	78	422	164.18	34.13
(%)	(1.2)	(7.0)	(18.0)	(20.1)	(25.9)	(13.7)	(7.0)	(5.2)	(1.5)	(0.6)	(100.0)				
急速進行性糸球体腎炎	5	12	31	57	41	34	26	10	7	4	227	57	284	168.74	39.65
(%)	(2.2)	(5.3)	(13.7)	(25.1)	(18.1)	(15.0)	(11.5)	(4.4)	(3.1)	(1.8)	(100.0)				
妊娠腎/妊娠中毒症	4	10	31	53	64	51	46	20	9	3	291	68	359	175.61	35.56
(%)	(1.4)	(3.4)	(10.7)	(18.2)	(22.0)	(17.5)	(15.8)	(6.9)	(3.1)	(1.0)	(100.0)				
その他の分類不能の腎炎	8	14	22	27	32	27	14	6			150	45	195	159.93	35.39
(%)	(5.3)	(9.3)	(14.7)	(18.0)	(21.3)	(18.0)	(9.3)	(4.0)			(100.0)				
多発性嚢胞腎	32	91	213	269	237	146	82	32	11	6	1,119	345	1,464	158.48	33.81
(%)	(2.9)	(8.1)	(19.0)	(24.0)	(21.2)	(13.0)	(7.3)	(2.9)	(1.0)	(0.5)	(100.0)				
腎硬化症	68	199	436	524	443	270	152	60	19	15	2,186	700	2,886	157.01	34.53
(%)	(3.1)	(9.1)	(19.9)	(24.0)	(20.3)	(12.4)	(7.0)	(2.7)	(0.9)	(0.7)	(100.0)				
悪性高血圧	4	15	42	52	60	28	20	9	3	3	236	76	312	164.52	37.02
(%)	(1.7)	(6.4)	(17.8)	(22.0)	(25.4)	(11.9)	(8.5)	(3.8)	(1.3)	(1.3)	(100.0)				
糖尿病性腎症	491	1,320	2,113	2,315	1,790	1,021	560	205	92	46	9,953	3,220	13,173	151.04	34.86
(%)	(4.9)	(13.3)	(21.2)	(23.3)	(18.0)	(10.3)	(5.6)	(2.1)	(0.9)	(0.5)	(100.0)				
SLE腎炎	5	17	43	44	56	45	34	12	12	8	276	87	363	171.95	41.08
(%)	(1.8)	(6.2)	(15.6)	(15.9)	(20.3)	(16.3)	(12.3)	(4.3)	(4.3)	(2.9)	(100.0)				
アミロイド腎	3	3	7	10	7	4	3	5	2		44	15	59	163.00	43.68
(%)	(6.8)	(6.8)	(15.9)	(22.7)	(15.9)	(9.1)	(6.8)	(11.4)	(4.5)		(100.0)				
痛風腎	6	12	22	30	24	10	7	2	4		117	37	154	154.13	35.30
(%)	(5.1)	(10.3)	(18.8)	(25.6)	(20.5)	(8.5)	(6.0)	(1.7)	(3.4)		(100.0)				
先天性代謝異常による腎不全		4	3	8	8	3	4	2		1	33	11	44	167.88	38.93
(%)		(12.1)	(9.1)	(24.2)	(24.2)	(9.1)	(12.1)	(6.1)		(3.0)	(100.0)				
腎・尿路結核		2		6	2	2	2	1		1	16	6	22	171.31	41.58
(%)		(12.5)		(37.5)	(12.5)	(12.5)	(12.5)	(6.3)		(6.3)	(100.0)				
腎・尿路結石	1	6	10	18	9	4	3	2		1	54	15	69	155.67	34.91
(%)	(1.9)	(11.1)	(18.5)	(33.3)	(16.7)	(7.4)	(5.6)	(3.7)		(1.9)	(100.0)				
腎・尿路腫瘍	3	5	17	18	15	5	2	3			68	17	85	150.96	31.95
(%)	(4.4)	(7.4)	(25.0)	(26.5)	(22.1)	(7.4)	(2.9)	(4.4)			(100.0)				
閉塞性尿路障害		4	7	6	18	6	3	5			49	25	74	166.49	32.92
(%)		(8.2)	(14.3)	(12.2)	(36.7)	(12.2)	(6.1)	(10.2)			(100.0)				
骨髄腫	1	1	4	6	4				1		17	2	19	147.71	34.88
(%)	(5.9)	(5.9)	(23.5)	(35.3)	(23.5)				(5.9)		(100.0)				
腎形成不全	1	7	17	12	18	12	2	3	3	1	76	25	101	161.00	37.23
(%)	(1.3)	(9.2)	(22.4)	(15.8)	(23.7)	(15.8)	(2.6)	(3.9)	(3.9)	(1.3)	(100.0)				
不明	58	256	471	559	527	365	197	78	40	25	2,576	904	3,480	160.45	36.21
(%)	(2.3)	(9.9)	(18.3)	(21.7)	(20.5)	(14.2)	(7.6)	(3.0)	(1.6)	(1.0)	(100.0)				
再導入	5	21	66	80	66	34	27	8	6	6	319	97	416	161.17	37.39
(%)	(1.6)	(6.6)	(20.7)	(25.1)	(20.7)	(10.7)	(8.5)	(2.5)	(1.9)	(1.9)	(100.0)				
その他	23	60	102	160	131	109	66	19	9	5	684	205	889	161.21	36.02
(%)	(3.4)	(8.8)	(14.9)	(23.4)	(19.2)	(15.9)	(9.6)	(2.8)	(1.3)	(0.7)	(100.0)				
合計	992	2,988	5,694	6,905	6,011	3,873	2,253	956	384	216	30,272	9,674	39,946	158.00	35.70
(%)	(3.3)	(9.9)	(18.8)	(22.8)	(19.9)	(12.8)	(7.4)	(3.2)	(1.3)	(0.7)	(100.0)				
記載なし															
(%)															
総計	992	2,988	5,694	6,905	6,011	3,873	2,253	956	384	216	30,272	9,674	39,946	158.00	35.70
(%)	(3.3)	(9.9)	(18.8)	(22.8)	(19.9)	(12.8)	(7.4)	(3.2)	(1.3)	(0.7)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査