

表 3704 総コレステロール濃度(mg/dL) Non-HDL-C濃度(mg/dL)別（腹膜透析患者）

Non-HDL-C濃度	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	合計	記載なし	総計	平均	標準偏差
50未満	9	6	1	1			1				18		18	105.06	33.58
(%)	(50.0)	(33.3)	(5.6)	(5.6)			(5.6)				(100.0)				
50～	25	46	22	13	5	1	2				114		114	117.99	24.54
(%)	(21.9)	(40.4)	(19.3)	(11.4)	(4.4)	(0.9)	(1.8)				(100.0)				
70～	4	69	147	85	27	13	5			1	351		351	136.83	22.20
(%)	(1.1)	(19.7)	(41.9)	(24.2)	(7.7)	(3.7)	(1.4)			(0.3)	(100.0)				
90～		12	169	324	126	23	4	1		1	660		660	149.87	16.83
(%)		(1.8)	(25.6)	(49.1)	(19.1)	(3.5)	(0.6)	(0.2)		(0.2)	(100.0)				
110～			14	223	354	171	25	8	1	1	797		797	169.66	17.27
(%)			(1.8)	(28.0)	(44.4)	(21.5)	(3.1)	(1.0)	(0.1)	(0.1)	(100.0)				
130～				11	181	315	126	27	1	1	662		662	188.77	16.33
(%)				(1.7)	(27.3)	(47.6)	(19.0)	(4.1)	(0.2)	(0.2)	(100.0)				
150～					8	153	179	74	17	5	436		436	207.54	17.89
(%)					(1.8)	(35.1)	(41.1)	(17.0)	(3.9)	(1.1)	(100.0)				
170～						8	112	102	30	11	263		263	224.37	16.21
(%)						(3.0)	(42.6)	(38.8)	(11.4)	(4.2)	(100.0)				
190～							1	49	44	26	120		120	247.37	18.54
(%)							(0.8)	(40.8)	(36.7)	(21.7)	(100.0)				
210～								1	20	69	90		90	287.54	34.73
(%)								(1.1)	(22.2)	(76.7)	(100.0)				
合計	38	133	353	657	701	684	455	262	113	115	3,511		3,511	178.73	39.85
(%)	(1.1)	(3.8)	(10.1)	(18.7)	(20.0)	(19.5)	(13.0)	(7.5)	(3.2)	(3.3)	(100.0)				
記載なし	12	27	95	149	164	149	109	56	46	34	841	4,589	5,430	180.17	43.30
(%)	(1.4)	(3.2)	(11.3)	(17.7)	(19.5)	(17.7)	(13.0)	(6.7)	(5.5)	(4.0)	(100.0)				
総計	50	160	448	806	865	833	564	318	159	149	4,352	4,589	8,941	179.01	40.54
(%)	(1.1)	(3.7)	(10.3)	(18.5)	(19.9)	(19.1)	(13.0)	(7.3)	(3.7)	(3.4)	(100.0)				
平均	56.00	71.77	89.20	103.60	119.69	137.11	154.22	171.55	189.70	222.58	128.47		128.47		
標準偏差	14.75	12.99	12.37	14.21	15.73	17.21	20.75	20.18	20.39	43.76	38.01		38.01		

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査