

表 3710 HDL-C濃度(mg/dL) 併用の状況別 (透析患者全体)

併用の状況	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	786	15,315	44,859	50,307	36,677	21,687	10,724	5,084	2,267	1,943	189,649	111,145	300,794	48.71	16.68
(%)	(0.4)	(8.1)	(23.7)	(26.5)	(19.3)	(11.4)	(5.7)	(2.7)	(1.2)	(1.0)	(100.0)				
PD単独	6	172	671	760	659	418	223	111	54	46	3,120	4,068	7,188	51.51	17.78
(%)	(0.2)	(5.5)	(21.5)	(24.4)	(21.1)	(13.4)	(7.1)	(3.6)	(1.7)	(1.5)	(100.0)				
HD(F)単独.PDカテあり		15	45	30	33	20	4	2	2	1	152	117	269	47.35	16.78
(%)		(9.9)	(29.6)	(19.7)	(21.7)	(13.2)	(2.6)	(1.3)	(1.3)	(0.7)	(100.0)				
PD.HD併用週1 HD(F)	1	108	230	202	157	81	29	23	6	10	847	697	1,544	46.73	17.18
(%)	(0.1)	(12.8)	(27.2)	(23.8)	(18.5)	(9.6)	(3.4)	(2.7)	(0.7)	(1.2)	(100.0)				
PD.HD併用週2 HD(F)	1	4	28	28	20	15	6	6	2		110	67	177	50.63	15.94
(%)	(0.9)	(3.6)	(25.5)	(25.5)	(18.2)	(13.6)	(5.5)	(5.5)	(1.8)		(100.0)				
PD.HD併用週3 HD(F)	1	4	2	9	5	6		1			28	12	40	46.96	15.35
(%)	(3.6)	(14.3)	(7.1)	(32.1)	(17.9)	(21.4)		(3.6)			(100.0)				
PD.HD併用週4 HD(F)												1	1		
(%)															
PD.HD併用その他	1	4	8	13	9	8	4	2		1	50	23	73	49.88	17.03
(%)	(2.0)	(8.0)	(16.0)	(26.0)	(18.0)	(16.0)	(8.0)	(4.0)		(2.0)	(100.0)				
合計	796	15,622	45,843	51,349	37,560	22,235	10,990	5,229	2,331	2,001	193,956	116,130	310,086	48.75	16.71
(%)	(0.4)	(8.1)	(23.6)	(26.5)	(19.4)	(11.5)	(5.7)	(2.7)	(1.2)	(1.0)	(100.0)				
不明		1	4	3	1	1	2				12	10	22	45.42	16.98
(%)		(8.3)	(33.3)	(25.0)	(8.3)	(8.3)	(16.7)				(100.0)				
記載なし															
(%)															
総計	796	15,623	45,847	51,352	37,561	22,236	10,992	5,229	2,331	2,001	193,968	116,140	310,108	48.75	16.71
(%)	(0.4)	(8.1)	(23.6)	(26.5)	(19.4)	(11.5)	(5.7)	(2.7)	(1.2)	(1.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査