

表 3714 HDL-C濃度(mg/dL) 原疾患別（透析患者全体）

原疾患	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	182	3,369	12,095	15,805	13,050	8,376	4,385	2,082	923	795	61,062	35,908	96,970	51.41	17.10
(%)	(0.3)	(5.5)	(19.8)	(25.9)	(21.4)	(13.7)	(7.2)	(3.4)	(1.5)	(1.3)	(100.0)				
慢性腎盂腎炎	6	102	330	466	396	268	145	82	30	39	1,864	1,178	3,042	52.95	18.18
(%)	(0.3)	(5.5)	(17.7)	(25.0)	(21.2)	(14.4)	(7.8)	(4.4)	(1.6)	(2.1)	(100.0)				
急速進行性糸球体腎炎	8	53	249	356	330	261	158	77	41	39	1,572	922	2,494	55.58	18.51
(%)	(0.5)	(3.4)	(15.8)	(22.6)	(21.0)	(16.6)	(10.1)	(4.9)	(2.6)	(2.5)	(100.0)				
妊娠腎/妊娠中毒症	1	32	147	255	245	171	115	51	30	13	1,060	554	1,614	55.42	17.08
(%)	(0.1)	(3.0)	(13.9)	(24.1)	(23.1)	(16.1)	(10.8)	(4.8)	(2.8)	(1.2)	(100.0)				
その他の分類不能の腎炎		40	155	224	198	128	71	37	14	10	877	496	1,373	52.84	16.73
(%)		(4.6)	(17.7)	(25.5)	(22.6)	(14.6)	(8.1)	(4.2)	(1.6)	(1.1)	(100.0)				
多発性嚢胞腎	22	361	1,427	1,946	1,585	955	464	197	94	81	7,132	3,874	11,006	50.85	16.46
(%)	(0.3)	(5.1)	(20.0)	(27.3)	(22.2)	(13.4)	(6.5)	(2.8)	(1.3)	(1.1)	(100.0)				
腎硬化症	52	1,298	4,178	5,002	3,641	2,069	959	446	191	149	17,985	10,313	28,298	48.67	16.00
(%)	(0.3)	(7.2)	(23.2)	(27.8)	(20.2)	(11.5)	(5.3)	(2.5)	(1.1)	(0.8)	(100.0)				
悪性高血圧	4	91	330	416	329	199	97	46	25	20	1,557	1,016	2,573	50.66	17.40
(%)	(0.3)	(5.8)	(21.2)	(26.7)	(21.1)	(12.8)	(6.2)	(3.0)	(1.6)	(1.3)	(100.0)				
糖尿病性腎症	404	8,504	21,220	19,840	12,294	6,339	2,802	1,316	572	498	73,789	44,292	118,081	45.27	15.56
(%)	(0.5)	(11.5)	(28.8)	(26.9)	(16.7)	(8.6)	(3.8)	(1.8)	(0.8)	(0.7)	(100.0)				
SLE腎炎	2	52	193	300	320	254	159	66	34	26	1,406	830	2,236	55.94	17.67
(%)	(0.1)	(3.7)	(13.7)	(21.3)	(22.8)	(18.1)	(11.3)	(4.7)	(2.4)	(1.8)	(100.0)				
アミロイド腎		8	42	70	64	48	20	19	7	10	288	163	451	56.53	20.06
(%)		(2.8)	(14.6)	(24.3)	(22.2)	(16.7)	(6.9)	(6.6)	(2.4)	(3.5)	(100.0)				
痛風腎	3	51	165	196	131	85	50	20	8	11	720	393	1,113	49.89	17.92
(%)	(0.4)	(7.1)	(22.9)	(27.2)	(18.2)	(11.8)	(6.9)	(2.8)	(1.1)	(1.5)	(100.0)				
先天性代謝異常による腎不全	1	10	22	37	41	18	11	8	3	5	156	116	272	53.69	19.79
(%)	(0.6)	(6.4)	(14.1)	(23.7)	(26.3)	(11.5)	(7.1)	(5.1)	(1.9)	(3.2)	(100.0)				
腎・尿路結核		8	26	37	23	20	6	6	2	3	131	75	206	51.59	17.77
(%)		(6.1)	(19.8)	(28.2)	(17.6)	(15.3)	(4.6)	(4.6)	(1.5)	(2.3)	(100.0)				
腎・尿路結石	2	19	61	108	80	52	19	14	2	5	362	219	581	51.40	16.66
(%)	(0.6)	(5.2)	(16.9)	(29.8)	(22.1)	(14.4)	(5.2)	(3.9)	(0.6)	(1.4)	(100.0)				
腎・尿路腫瘍	3	52	127	163	117	75	27	14	2	7	587	299	886	48.30	17.16
(%)	(0.5)	(8.9)	(21.6)	(27.8)	(19.9)	(12.8)	(4.6)	(2.4)	(0.3)	(1.2)	(100.0)				
閉塞性尿路障害	2	19	93	123	97	74	30	8	11	7	464	259	723	51.63	17.05
(%)	(0.4)	(4.1)	(20.0)	(26.5)	(20.9)	(15.9)	(6.5)	(1.7)	(2.4)	(1.5)	(100.0)				
骨髄腫	1	9	38	42	35	24	8	3	4	2	166	110	276	49.87	16.93
(%)	(0.6)	(5.4)	(22.9)	(25.3)	(21.1)	(14.5)	(4.8)	(1.8)	(2.4)	(1.2)	(100.0)				
腎形成不全	2	20	73	111	94	51	37	21	6	7	422	227	649	52.50	17.27
(%)	(0.5)	(4.7)	(17.3)	(26.3)	(22.3)	(12.1)	(8.8)	(5.0)	(1.4)	(1.7)	(100.0)				
不明	70	1,130	3,656	4,354	3,310	1,982	1,011	498	219	174	16,404	11,140	27,544	49.68	16.81
(%)	(0.4)	(6.9)	(22.3)	(26.5)	(20.2)	(12.1)	(6.2)	(3.0)	(1.3)	(1.1)	(100.0)				
再導入	4	92	259	354	300	180	106	62	27	32	1,416	747	2,163	52.32	18.15
(%)	(0.3)	(6.5)	(18.3)	(25.0)	(21.2)	(12.7)	(7.5)	(4.4)	(1.9)	(2.3)	(100.0)				
その他	27	303	961	1,147	881	607	312	156	86	68	4,548	3,009	7,557	50.96	17.83
(%)	(0.6)	(6.7)	(21.1)	(25.2)	(19.4)	(13.3)	(6.9)	(3.4)	(1.9)	(1.5)	(100.0)				
合計	796	15,623	45,847	51,352	37,561	22,236	10,992	5,229	2,331	2,001	193,968	116,140	310,108	48.75	16.71
(%)	(0.4)	(8.1)	(23.6)	(26.5)	(19.4)	(11.5)	(5.7)	(2.7)	(1.2)	(1.0)	(100.0)				
記載なし															
(%)															
総計	796	15,623	45,847	51,352	37,561	22,236	10,992	5,229	2,331	2,001	193,968	116,140	310,108	48.75	16.71
(%)	(0.4)	(8.1)	(23.6)	(26.5)	(19.4)	(11.5)	(5.7)	(2.7)	(1.2)	(1.0)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査