

表 3716 HDL-C濃度(mg/dL) 週透析回数(回/週)別 (透析患者全体)

週透析回数	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	合計	記載なし	総計	平均	標準偏差
1回	4	132	306	297	227	138	53	34	10	17	1,218	622	1,840	47.93	17.52
(%)	(0.3)	(10.8)	(25.1)	(24.4)	(18.6)	(11.3)	(4.4)	(2.8)	(0.8)	(1.4)	(100.0)				
2回	21	289	968	1,068	812	488	229	108	44	53	4,080	2,112	6,192	49.16	16.81
(%)	(0.5)	(7.1)	(23.7)	(26.2)	(19.9)	(12.0)	(5.6)	(2.6)	(1.1)	(1.3)	(100.0)				
3回	755	14,789	43,218	48,405	35,300	20,829	10,315	4,878	2,179	1,850	182,518	88,121	270,639	48.69	16.67
(%)	(0.4)	(8.1)	(23.7)	(26.5)	(19.3)	(11.4)	(5.7)	(2.7)	(1.2)	(1.0)	(100.0)				
4回	3	52	151	190	117	93	51	22	11	14	704	290	994	50.44	18.09
(%)	(0.4)	(7.4)	(21.4)	(27.0)	(16.6)	(13.2)	(7.2)	(3.1)	(1.6)	(2.0)	(100.0)				
5回		3	5	11	10	11	7	4	3	3	57	32	89	60.79	21.38
(%)		(5.3)	(8.8)	(19.3)	(17.5)	(19.3)	(12.3)	(7.0)	(5.3)	(5.3)	(100.0)				
6回		3	4	7	7	9	6	4	1		41	40	81	57.51	17.64
(%)		(7.3)	(9.8)	(17.1)	(17.1)	(22.0)	(14.6)	(9.8)	(2.4)		(100.0)				
7回		1	4	2	1	4	2	2	2	4	22	8	30	70.77	30.41
(%)		(4.5)	(18.2)	(9.1)	(4.5)	(18.2)	(9.1)	(9.1)	(9.1)	(18.2)	(100.0)				
合計	783	15,269	44,656	49,980	36,474	21,572	10,663	5,052	2,250	1,941	188,640	91,225	279,865	48.71	16.69
(%)	(0.4)	(8.1)	(23.7)	(26.5)	(19.3)	(11.4)	(5.7)	(2.7)	(1.2)	(1.0)	(100.0)				
記載なし	13	354	1,191	1,372	1,087	664	329	177	81	60	5,328	24,915	30,243	50.18	17.12
(%)	(0.2)	(6.6)	(22.4)	(25.8)	(20.4)	(12.5)	(6.2)	(3.3)	(1.5)	(1.1)	(100.0)				
総計	796	15,623	45,847	51,352	37,561	22,236	10,992	5,229	2,331	2,001	193,968	116,140	310,108	48.75	16.71
(%)	(0.4)	(8.1)	(23.6)	(26.5)	(19.4)	(11.5)	(5.7)	(2.7)	(1.2)	(1.0)	(100.0)				
平均	2.97	2.97	2.97	2.97	2.97	2.97	2.98	2.98	2.98	2.97	2.97	2.97	2.97		
標準偏差	0.22	0.24	0.23	0.23	0.23	0.25	0.24	0.26	0.26	0.33	0.24	0.24	0.24		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査