

表 3734 HDL-C濃度(mg/dL) 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	19	469	1,822	2,651	2,411	1,730	925	488	194	190	10,899	4,177	15,076	53.73	17.50
(%)	(0.2)	(4.3)	(16.7)	(24.3)	(22.1)	(15.9)	(8.5)	(4.5)	(1.8)	(1.7)	(100.0)				
慢性腎盂腎炎	1	8	49	82	68	55	23	16	5	7	314	108	422	54.71	17.42
(%)	(0.3)	(2.5)	(15.6)	(26.1)	(21.7)	(17.5)	(7.3)	(5.1)	(1.6)	(2.2)	(100.0)				
急速進行性糸球体腎炎		4	32	44	40	34	31	15	7	7	214	70	284	58.49	19.07
(%)		(1.9)	(15.0)	(20.6)	(18.7)	(15.9)	(14.5)	(7.0)	(3.3)	(3.3)	(100.0)				
妊娠腎/妊娠中毒症		6	30	51	63	54	39	18	12	4	277	82	359	59.15	17.38
(%)		(2.2)	(10.8)	(18.4)	(22.7)	(19.5)	(14.1)	(6.5)	(4.3)	(1.4)	(100.0)				
その他の分類不能の腎炎		10	23	33	26	19	18	6	2	3	140	55	195	53.96	18.50
(%)		(7.1)	(16.4)	(23.6)	(18.6)	(13.6)	(12.9)	(4.3)	(1.4)	(2.1)	(100.0)				
多発性嚢胞腎	2	41	204	273	248	164	82	26	19	10	1,069	395	1,464	52.08	16.03
(%)	(0.2)	(3.8)	(19.1)	(25.5)	(23.2)	(15.3)	(7.7)	(2.4)	(1.8)	(0.9)	(100.0)				
腎硬化症	2	147	464	569	471	284	123	60	25	25	2,170	716	2,886	49.94	16.17
(%)	(0.1)	(6.8)	(21.4)	(26.2)	(21.7)	(13.1)	(5.7)	(2.8)	(1.2)	(1.2)	(100.0)				
悪性高血圧		11	34	65	51	34	16	10	6	3	230	82	312	53.50	18.00
(%)		(4.8)	(14.8)	(28.3)	(22.2)	(14.8)	(7.0)	(4.3)	(2.6)	(1.3)	(100.0)				
糖尿病性腎症	46	982	2,752	2,545	1,692	851	415	197	76	60	9,616	3,557	13,173	45.95	15.62
(%)	(0.5)	(10.2)	(28.6)	(26.5)	(17.6)	(8.8)	(4.3)	(2.0)	(0.8)	(0.6)	(100.0)				
SLE腎炎		4	22	57	62	52	42	14	7	8	268	95	363	59.79	17.14
(%)		(1.5)	(8.2)	(21.3)	(23.1)	(19.4)	(15.7)	(5.2)	(2.6)	(3.0)	(100.0)				
アミロイド腎		1	10	10	9	7	1	6	2	2	48	11	59	57.08	21.53
(%)		(2.1)	(20.8)	(20.8)	(18.8)	(14.6)	(2.1)	(12.5)	(4.2)	(4.2)	(100.0)				
痛風腎		7	25	30	18	17	8	4	1	1	111	43	154	50.40	16.37
(%)		(6.3)	(22.5)	(27.0)	(16.2)	(15.3)	(7.2)	(3.6)	(0.9)	(0.9)	(100.0)				
先天性代謝異常による腎不全			2	5	7	5	5	2	2	1	29	15	44	62.14	18.90
(%)			(6.9)	(17.2)	(24.1)	(17.2)	(17.2)	(6.9)	(6.9)	(3.4)	(100.0)				
腎・尿路結核			2	6	3	3	1	1		1	17	5	22	57.47	19.28
(%)			(11.8)	(35.3)	(17.6)	(17.6)	(5.9)	(5.9)		(5.9)	(100.0)				
腎・尿路結石			7	13	12	6	5	1			44	25	69	52.55	13.66
(%)			(15.9)	(29.5)	(27.3)	(13.6)	(11.4)	(2.3)			(100.0)				
腎・尿路腫瘍	1	3	8	23	17	15	3	2		1	73	12	85	52.07	15.89
(%)	(1.4)	(4.1)	(11.0)	(31.5)	(23.3)	(20.5)	(4.1)	(2.7)		(1.4)	(100.0)				
閉塞性尿路障害		2	8	15	14	7	3	3	2		54	20	74	53.07	15.65
(%)		(3.7)	(14.8)	(27.8)	(25.9)	(13.0)	(5.6)	(5.6)	(3.7)		(100.0)				
骨髄腫			2	6	7	2					17	2	19	50.12	7.56
(%)			(11.8)	(35.3)	(41.2)	(11.8)					(100.0)				
腎形成不全		3	8	17	22	11	8	5		2	76	25	101	55.64	16.44
(%)		(3.9)	(10.5)	(22.4)	(28.9)	(14.5)	(10.5)	(6.6)		(2.6)	(100.0)				
不明	7	136	440	622	526	370	186	106	44	34	2,471	1,009	3,480	52.43	17.39
(%)	(0.3)	(5.5)	(17.8)	(25.2)	(21.3)	(15.0)	(7.5)	(4.3)	(1.8)	(1.4)	(100.0)				
再導入	1	17	57	70	73	39	26	12	4	11	310	106	416	53.63	19.16
(%)	(0.3)	(5.5)	(18.4)	(22.6)	(23.5)	(12.6)	(8.4)	(3.9)	(1.3)	(3.5)	(100.0)				
その他	2	23	130	173	131	98	63	21	23	13	677	212	889	53.62	18.10
(%)	(0.3)	(3.4)	(19.2)	(25.6)	(19.4)	(14.5)	(9.3)	(3.1)	(3.4)	(1.9)	(100.0)				
合計	81	1,874	6,131	7,360	5,971	3,857	2,023	1,013	431	383	29,124	10,822	39,946	50.85	17.18
(%)	(0.3)	(6.4)	(21.1)	(25.3)	(20.5)	(13.2)	(6.9)	(3.5)	(1.5)	(1.3)	(100.0)				
記載なし															
(%)															
総計	81	1,874	6,131	7,360	5,971	3,857	2,023	1,013	431	383	29,124	10,822	39,946	50.85	17.18
(%)	(0.3)	(6.4)	(21.1)	(25.3)	(20.5)	(13.2)	(6.9)	(3.5)	(1.5)	(1.3)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査