

表 3736 HDL-C濃度(mg/dL) 透析後体重(kg)別（血液透析濾過患者 透析回数週3回）

透析後体重	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	合計	記載なし	総計	平均	標準偏差
20未満			2	2		1	2				7	5	12	54.43	17.21
(%)			(28.6)	(28.6)		(14.3)	(28.6)				(100.0)				
20～			7	11	15	9	9	3			54	22	76	56.17	14.56
(%)			(13.0)	(20.4)	(27.8)	(16.7)	(16.7)	(5.6)			(100.0)				
30～	4	55	226	392	441	340	236	118	59	64	1,935	747	2,682	58.26	19.00
(%)	(0.2)	(2.8)	(11.7)	(20.3)	(22.8)	(17.6)	(12.2)	(6.1)	(3.0)	(3.3)	(100.0)				
40～	18	255	1,006	1,652	1,652	1,351	721	399	169	134	7,357	2,648	10,005	55.79	17.45
(%)	(0.2)	(3.5)	(13.7)	(22.5)	(22.5)	(18.4)	(9.8)	(5.4)	(2.3)	(1.8)	(100.0)				
50～	30	449	1,767	2,446	2,073	1,319	647	324	134	112	9,301	3,189	12,490	51.68	16.67
(%)	(0.3)	(4.8)	(19.0)	(26.3)	(22.3)	(14.2)	(7.0)	(3.5)	(1.4)	(1.2)	(100.0)				
60～	14	486	1,665	1,733	1,218	605	295	131	57	59	6,263	2,102	8,365	47.51	15.91
(%)	(0.2)	(7.8)	(26.6)	(27.7)	(19.4)	(9.7)	(4.7)	(2.1)	(0.9)	(0.9)	(100.0)				
70～	7	320	813	737	395	166	73	30	9	9	2,559	864	3,423	43.34	13.85
(%)	(0.3)	(12.5)	(31.8)	(28.8)	(15.4)	(6.5)	(2.9)	(1.2)	(0.4)	(0.4)	(100.0)				
80～	2	168	373	250	113	44	27	5	2	2	986	287	1,273	40.49	12.79
(%)	(0.2)	(17.0)	(37.8)	(25.4)	(11.5)	(4.5)	(2.7)	(0.5)	(0.2)	(0.2)	(100.0)				
90～	4	81	163	69	34	7	6	2		1	367	88	455	37.27	11.83
(%)	(1.1)	(22.1)	(44.4)	(18.8)	(9.3)	(1.9)	(1.6)	(0.5)		(0.3)	(100.0)				
100～	2	27	47	19	5	3	3				106	36	142	36.25	11.49
(%)	(1.9)	(25.5)	(44.3)	(17.9)	(4.7)	(2.8)	(2.8)				(100.0)				
110～		16	15	9	5	2					47	14	61	35.81	11.05
(%)		(34.0)	(31.9)	(19.1)	(10.6)	(4.3)					(100.0)				
120～		10	14	7	1						32	5	37	33.75	8.02
(%)		(31.3)	(43.8)	(21.9)	(3.1)						(100.0)				
合計	81	1,867	6,098	7,327	5,952	3,847	2,019	1,012	430	381	29,014	10,007	39,021	50.87	17.18
(%)	(0.3)	(6.4)	(21.0)	(25.3)	(20.5)	(13.3)	(7.0)	(3.5)	(1.5)	(1.3)	(100.0)				
記載なし		7	33	33	19	10	4	1	1	2	110	815	925	46.96	16.24
(%)		(6.4)	(30.0)	(30.0)	(17.3)	(9.1)	(3.6)	(0.9)	(0.9)	(1.8)	(100.0)				
総計	81	1,874	6,131	7,360	5,971	3,857	2,023	1,013	431	383	29,124	10,822	39,946	50.85	17.18
(%)	(0.3)	(6.4)	(21.1)	(25.3)	(20.5)	(13.2)	(6.9)	(3.5)	(1.5)	(1.3)	(100.0)				
平均	59.58	65.13	61.51	57.61	54.86	52.57	51.61	50.79	49.97	50.24	56.82	56.06	56.63		
標準偏差	15.83	15.99	14.27	12.56	11.41	10.49	10.75	9.78	9.79	10.14	13.12	12.77	13.03		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査