

表 3743 HDL-C濃度(mg/dL) 透析歴(年)別 (腹膜透析患者)

透析歴	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	合計	記載なし	総計	平均	標準偏差
2年未満	1	82	348	390	314	205	118	72	24	29	1,583	2,017	3,600	52.15	18.79
(%)	(0.1)	(5.2)	(22.0)	(24.6)	(19.8)	(13.0)	(7.5)	(4.5)	(1.5)	(1.8)	(100.0)				
2年～	5	117	323	346	294	194	79	35	24	17	1,434	1,767	3,201	49.64	17.14
(%)	(0.3)	(8.2)	(22.5)	(24.1)	(20.5)	(13.5)	(5.5)	(2.4)	(1.7)	(1.2)	(100.0)				
5年～	1	72	204	179	168	91	54	30	12	9	820	837	1,657	49.59	17.36
(%)	(0.1)	(8.8)	(24.9)	(21.8)	(20.5)	(11.1)	(6.6)	(3.7)	(1.5)	(1.1)	(100.0)				
10年～	1	9	39	52	42	19	4	3	1	1	171	169	340	47.81	13.89
(%)	(0.6)	(5.3)	(22.8)	(30.4)	(24.6)	(11.1)	(2.3)	(1.8)	(0.6)	(0.6)	(100.0)				
15年～		3	12	17	15	6	2			1	56	41	97	47.98	13.63
(%)		(5.4)	(21.4)	(30.4)	(26.8)	(10.7)	(3.6)			(1.8)	(100.0)				
20年～		1	2	3	2	1	2				11	9	20	51.18	16.95
(%)		(9.1)	(18.2)	(27.3)	(18.2)	(9.1)	(18.2)				(100.0)				
25年～		2	1	2	3						8		8	42.88	13.20
(%)		(25.0)	(12.5)	(25.0)	(37.5)						(100.0)				
30年～		2	3	1	1						7	3	10	37.14	10.53
(%)		(28.6)	(42.9)	(14.3)	(14.3)						(100.0)				
35年～			1					1			2	1	3	61.00	36.77
(%)			(50.0)					(50.0)			(100.0)				
40年～															
(%)															
合計	8	288	933	990	839	516	259	141	61	57	4,092	4,844	8,936	50.47	17.72
(%)	(0.2)	(7.0)	(22.8)	(24.2)	(20.5)	(12.6)	(6.3)	(3.4)	(1.5)	(1.4)	(100.0)				
不明												5	5		
(%)															
記載なし															
(%)															
総計	8	288	933	990	839	516	259	141	61	57	4,092	4,849	8,941	50.47	17.72
(%)	(0.2)	(7.0)	(22.8)	(24.2)	(20.5)	(12.6)	(6.3)	(3.4)	(1.5)	(1.4)	(100.0)				
平均	4.13	3.87	3.46	3.35	3.45	3.04	2.82	2.75	2.64	2.47	3.32	2.91	3.10		
標準偏差	4.12	4.35	4.06	3.87	3.93	3.25	3.35	4.03	2.53	2.95	3.85	3.30	3.56		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査