

表 3750 HDL-C濃度(mg/dL) 脈拍(拍/分)別 (腹膜透析患者)

脈拍	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	合計	記載なし	総計	平均	標準偏差
40未満				1	2						3	2	5	51.00	7.00
(%)				(33.3)	(66.7)						(100.0)				
40～		1	6	2	4	2	2				17	8	25	48.88	14.25
(%)		(5.9)	(35.3)	(11.8)	(23.5)	(11.8)	(11.8)				(100.0)				
50～		17	29	49	21	13	4	1			134	53	187	44.28	12.93
(%)		(12.7)	(21.6)	(36.6)	(15.7)	(9.7)	(3.0)	(0.7)			(100.0)				
60～	1	57	153	159	115	73	34	20	11	6	629	221	850	49.00	18.04
(%)	(0.2)	(9.1)	(24.3)	(25.3)	(18.3)	(11.6)	(5.4)	(3.2)	(1.7)	(1.0)	(100.0)				
70～	3	70	227	239	204	139	66	29	12	12	1,001	270	1,271	50.33	17.02
(%)	(0.3)	(7.0)	(22.7)	(23.9)	(20.4)	(13.9)	(6.6)	(2.9)	(1.2)	(1.2)	(100.0)				
80～	1	49	165	188	165	92	52	30	15	11	768	256	1,024	51.24	17.61
(%)	(0.1)	(6.4)	(21.5)	(24.5)	(21.5)	(12.0)	(6.8)	(3.9)	(2.0)	(1.4)	(100.0)				
90～	1	30	85	78	79	47	26	8	10	6	370	111	481	50.54	17.79
(%)	(0.3)	(8.1)	(23.0)	(21.1)	(21.4)	(12.7)	(7.0)	(2.2)	(2.7)	(1.6)	(100.0)				
100～		10	28	42	30	19	8	9	1	2	149	55	204	51.49	17.14
(%)		(6.7)	(18.8)	(28.2)	(20.1)	(12.8)	(5.4)	(6.0)	(0.7)	(1.3)	(100.0)				
110～		1	12	6	8	8	1	2			38	14	52	50.66	14.90
(%)		(2.6)	(31.6)	(15.8)	(21.1)	(21.1)	(2.6)	(5.3)			(100.0)				
120～	1	1	3			3					8	12	20	40.63	19.48
(%)	(12.5)	(12.5)	(37.5)			(37.5)					(100.0)				
合計	7	236	708	764	628	396	193	99	49	37	3,117	1,002	4,119	50.08	17.33
(%)	(0.2)	(7.6)	(22.7)	(24.5)	(20.1)	(12.7)	(6.2)	(3.2)	(1.6)	(1.2)	(100.0)				
記載なし	1	52	225	226	211	120	66	42	12	20	975	3,847	4,822	51.74	18.86
(%)	(0.1)	(5.3)	(23.1)	(23.2)	(21.6)	(12.3)	(6.8)	(4.3)	(1.2)	(2.1)	(100.0)				
総計	8	288	933	990	839	516	259	141	61	57	4,092	4,849	8,941	50.47	17.72
(%)	(0.2)	(7.0)	(22.8)	(24.2)	(20.5)	(12.6)	(6.3)	(3.4)	(1.5)	(1.4)	(100.0)				
平均	84.57	76.57	77.74	77.15	78.22	79.11	78.93	80.76	79.69	80.03	78.02	78.29	78.09		
標準偏差	20.53	12.86	13.15	13.07	12.67	13.34	11.73	12.56	11.31	11.03	12.92	14.29	13.27		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査