

表 3753 Non-HDL-C濃度(mg/dL) 併用の状況別（透析患者全体）

併用の状況	50未満	50～	70～	90～	110～	130～	150～	170～	190～	210～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	3,391	15,970	35,182	41,665	33,356	20,276	10,139	4,453	1,686	1,030	167,148	133,646	300,794	106.63	33.36
(%)	(2.0)	(9.6)	(21.0)	(24.9)	(20.0)	(12.1)	(6.1)	(2.7)	(1.0)	(0.6)	(100.0)				
PD単独	12	76	241	453	596	522	354	213	102	83	2,652	4,536	7,188	131.38	38.78
(%)	(0.5)	(2.9)	(9.1)	(17.1)	(22.5)	(19.7)	(13.3)	(8.0)	(3.8)	(3.1)	(100.0)				
HD(F)単独.PDカテあり	1	11	25	39	30	17	9	4	3	2	141	128	269	111.99	34.26
(%)	(0.7)	(7.8)	(17.7)	(27.7)	(21.3)	(12.1)	(6.4)	(2.8)	(2.1)	(1.4)	(100.0)				
PD.HD併用週1 HD(F)	6	31	96	183	171	124	67	42	13	7	740	804	1,544	118.94	33.78
(%)	(0.8)	(4.2)	(13.0)	(24.7)	(23.1)	(16.8)	(9.1)	(5.7)	(1.8)	(0.9)	(100.0)				
PD.HD併用週2 HD(F)		3	13	23	24	14	13	9	4	1	104	73	177	125.45	36.78
(%)		(2.9)	(12.5)	(22.1)	(23.1)	(13.5)	(12.5)	(8.7)	(3.8)	(1.0)	(100.0)				
PD.HD併用週3 HD(F)		2	5	4	7	4	4				26	14	40	114.08	29.75
(%)		(7.7)	(19.2)	(15.4)	(26.9)	(15.4)	(15.4)				(100.0)				
PD.HD併用週4 HD(F)												1	1		
(%)															
PD.HD併用その他		6	5	9	14	6	4	1	1		46	27	73	112.89	32.99
(%)		(13.0)	(10.9)	(19.6)	(30.4)	(13.0)	(8.7)	(2.2)	(2.2)		(100.0)				
合計	3,410	16,099	35,567	42,376	34,198	20,963	10,590	4,722	1,809	1,123	170,857	139,229	310,086	107.08	33.61
(%)	(2.0)	(9.4)	(20.8)	(24.8)	(20.0)	(12.3)	(6.2)	(2.8)	(1.1)	(0.7)	(100.0)				
不明		2	2	2	5					1	12	10	22	106.25	43.83
(%)		(16.7)	(16.7)	(16.7)	(41.7)					(8.3)	(100.0)				
記載なし (%)															
総計	3,410	16,101	35,569	42,378	34,203	20,963	10,590	4,722	1,809	1,124	170,869	139,239	310,108	107.08	33.61
(%)	(2.0)	(9.4)	(20.8)	(24.8)	(20.0)	(12.3)	(6.2)	(2.8)	(1.1)	(0.7)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査