

表 3755 Non-HDL-C濃度(mg/dL) 年齢(歳)別 (透析患者全体)

年齢	50未満	50～	70～	90～	110～	130～	150～	170～	190～	210～	合計	記載なし	総計	平均	標準偏差
15歳未満		2	5		2	3	3		2	2	19	98	117	135.37	61.83
(%)		(10.5)	(26.3)		(10.5)	(15.8)	(15.8)		(10.5)	(10.5)	(100.0)				
15歳～	18	77	124	140	92	39	28	9	3	2	532	505	1,037	99.08	31.70
(%)	(3.4)	(14.5)	(23.3)	(26.3)	(17.3)	(7.3)	(5.3)	(1.7)	(0.6)	(0.4)	(100.0)				
30歳～	130	654	1,574	1,965	1,673	1,077	579	250	106	77	8,085	6,300	14,385	110.19	34.72
(%)	(1.6)	(8.1)	(19.5)	(24.3)	(20.7)	(13.3)	(7.2)	(3.1)	(1.3)	(1.0)	(100.0)				
45歳～	597	2,713	5,846	7,395	6,449	4,417	2,405	1,152	451	286	31,711	24,857	56,568	110.96	35.34
(%)	(1.9)	(8.6)	(18.4)	(23.3)	(20.3)	(13.9)	(7.6)	(3.6)	(1.4)	(0.9)	(100.0)				
60歳～	1,563	7,456	16,321	19,191	15,848	9,719	4,950	2,185	841	509	78,583	63,395	141,978	107.20	33.65
(%)	(2.0)	(9.5)	(20.8)	(24.4)	(20.2)	(12.4)	(6.3)	(2.8)	(1.1)	(0.6)	(100.0)				
75歳～	1,050	4,932	11,096	12,988	9,643	5,425	2,491	1,069	383	232	49,309	41,688	90,997	104.13	31.93
(%)	(2.1)	(10.0)	(22.5)	(26.3)	(19.6)	(11.0)	(5.1)	(2.2)	(0.8)	(0.5)	(100.0)				
90歳～	52	267	603	698	496	283	134	57	23	16	2,629	2,376	5,005	104.03	32.13
(%)	(2.0)	(10.2)	(22.9)	(26.6)	(18.9)	(10.8)	(5.1)	(2.2)	(0.9)	(0.6)	(100.0)				
合計	3,410	16,101	35,569	42,377	34,203	20,963	10,590	4,722	1,809	1,124	170,868	139,219	310,087	107.08	33.61
(%)	(2.0)	(9.4)	(20.8)	(24.8)	(20.0)	(12.3)	(6.2)	(2.8)	(1.1)	(0.7)	(100.0)				
不明				1							1	20	21	102.00	
(%)				(100.0)							(100.0)				
記載なし															
(%)															
総計	3,410	16,101	35,569	42,378	34,203	20,963	10,590	4,722	1,809	1,124	170,869	139,239	310,108	107.08	33.61
(%)	(2.0)	(9.4)	(20.8)	(24.8)	(20.0)	(12.3)	(6.2)	(2.8)	(1.1)	(0.7)	(100.0)				
平均	67.97	68.10	68.21	67.88	67.16	66.40	65.60	65.26	64.92	64.25	67.38	67.74	67.54		
標準偏差	12.27	12.31	12.36	12.48	12.38	12.36	12.37	12.38	12.35	12.99	12.43	12.57	12.49		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査