

表 3757 Non-HDL-C濃度(mg/dL) 原疾患別（透析患者全体）

原疾患	50未満	50～	70～	90～	110～	130～	150～	170～	190～	210～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	991	4,614	10,760	13,313	11,222	6,956	3,595	1,550	553	324	53,878	43,092	96,970	108.35	33.22
(%)	(1.8)	(8.6)	(20.0)	(24.7)	(20.8)	(12.9)	(6.7)	(2.9)	(1.0)	(0.6)	(100.0)				
慢性腎盂腎炎	33	134	298	401	357	221	110	59	16	7	1,636	1,406	3,042	109.62	33.20
(%)	(2.0)	(8.2)	(18.2)	(24.5)	(21.8)	(13.5)	(6.7)	(3.6)	(1.0)	(0.4)	(100.0)				
急速進行性糸球体腎炎	19	76	235	344	278	184	126	63	19	27	1,371	1,123	2,494	115.55	36.67
(%)	(1.4)	(5.5)	(17.1)	(25.1)	(20.3)	(13.4)	(9.2)	(4.6)	(1.4)	(2.0)	(100.0)				
妊娠腎/妊娠中毒症	6	37	144	229	211	160	89	42	18	7	943	671	1,614	117.87	33.20
(%)	(0.6)	(3.9)	(15.3)	(24.3)	(22.4)	(17.0)	(9.4)	(4.5)	(1.9)	(0.7)	(100.0)				
その他の分類不能の腎炎	14	77	160	189	146	102	48	26	8	5	775	598	1,373	107.74	34.52
(%)	(1.8)	(9.9)	(20.6)	(24.4)	(18.8)	(13.2)	(6.2)	(3.4)	(1.0)	(0.6)	(100.0)				
多発性嚢胞腎	90	486	1,239	1,569	1,391	860	386	171	41	30	6,263	4,743	11,006	108.54	31.63
(%)	(1.4)	(7.8)	(19.8)	(25.1)	(22.2)	(13.7)	(6.2)	(2.7)	(0.7)	(0.5)	(100.0)				
腎硬化症	250	1,332	3,310	4,208	3,303	1,913	929	353	178	105	15,881	12,417	28,298	107.26	32.40
(%)	(1.6)	(8.4)	(20.8)	(26.5)	(20.8)	(12.0)	(5.8)	(2.2)	(1.1)	(0.7)	(100.0)				
悪性高血圧	22	94	277	334	290	178	84	51	20	13	1,363	1,210	2,573	110.80	35.84
(%)	(1.6)	(6.9)	(20.3)	(24.5)	(21.3)	(13.1)	(6.2)	(3.7)	(1.5)	(1.0)	(100.0)				
糖尿病性腎症	1,535	7,159	14,432	16,034	12,225	7,277	3,693	1,726	700	441	65,222	52,859	118,081	104.90	34.01
(%)	(2.4)	(11.0)	(22.1)	(24.6)	(18.7)	(11.2)	(5.7)	(2.6)	(1.1)	(0.7)	(100.0)				
SLE腎炎	23	86	227	285	247	186	120	42	26	16	1,258	978	2,236	114.52	37.51
(%)	(1.8)	(6.8)	(18.0)	(22.7)	(19.6)	(14.8)	(9.5)	(3.3)	(2.1)	(1.3)	(100.0)				
アミロイド腎	13	35	47	52	46	25	13	10	6	1	248	203	451	103.32	38.85
(%)	(5.2)	(14.1)	(19.0)	(21.0)	(18.5)	(10.1)	(5.2)	(4.0)	(2.4)	(0.4)	(100.0)				
痛風腎	16	65	128	151	134	75	36	21	9	2	637	476	1,113	106.57	33.91
(%)	(2.5)	(10.2)	(20.1)	(23.7)	(21.0)	(11.8)	(5.7)	(3.3)	(1.4)	(0.3)	(100.0)				
先天性代謝異常による腎不全	5	15	24	33	24	18	11	4	3	2	139	133	272	108.61	38.06
(%)	(3.6)	(10.8)	(17.3)	(23.7)	(17.3)	(12.9)	(7.9)	(2.9)	(2.2)	(1.4)	(100.0)				
腎・尿路結核	1	13	23	20	34	11	10	2		1	115	91	206	106.83	32.21
(%)	(0.9)	(11.3)	(20.0)	(17.4)	(29.6)	(9.6)	(8.7)	(1.7)		(0.9)	(100.0)				
腎・尿路結石	4	37	67	78	57	41	20	7	1		312	269	581	104.32	30.36
(%)	(1.3)	(11.9)	(21.5)	(25.0)	(18.3)	(13.1)	(6.4)	(2.2)	(0.3)		(100.0)				
腎・尿路腫瘍	10	49	126	129	93	55	34	15	7	2	520	366	886	105.75	33.38
(%)	(1.9)	(9.4)	(24.2)	(24.8)	(17.9)	(10.6)	(6.5)	(2.9)	(1.3)	(0.4)	(100.0)				
閉塞性尿路障害	6	32	75	89	89	62	24	17	4	4	402	321	723	111.65	35.24
(%)	(1.5)	(8.0)	(18.7)	(22.1)	(22.1)	(15.4)	(6.0)	(4.2)	(1.0)	(1.0)	(100.0)				
骨髄腫	1	17	23	36	26	19	14	7	2	5	150	126	276	115.08	39.71
(%)	(0.7)	(11.3)	(15.3)	(24.0)	(17.3)	(12.7)	(9.3)	(4.7)	(1.3)	(3.3)	(100.0)				
腎形成不全	8	28	65	84	77	46	32	9	4	2	355	294	649	110.74	33.43
(%)	(2.3)	(7.9)	(18.3)	(23.7)	(21.7)	(13.0)	(9.0)	(2.5)	(1.1)	(0.6)	(100.0)				
不明	274	1,238	2,894	3,553	2,883	1,878	874	413	133	86	14,226	13,318	27,544	107.80	33.24
(%)	(1.9)	(8.7)	(20.3)	(25.0)	(20.3)	(13.2)	(6.1)	(2.9)	(0.9)	(0.6)	(100.0)				
再導入	23	112	219	328	248	177	92	36	9	13	1,257	906	2,163	109.88	35.23
(%)	(1.8)	(8.9)	(17.4)	(26.1)	(19.7)	(14.1)	(7.3)	(2.9)	(0.7)	(1.0)	(100.0)				
その他	66	365	796	919	822	519	250	98	52	31	3,918	3,639	7,557	108.50	34.14
(%)	(1.7)	(9.3)	(20.3)	(23.5)	(21.0)	(13.2)	(6.4)	(2.5)	(1.3)	(0.8)	(100.0)				
合計	3,410	16,101	35,569	42,378	34,203	20,963	10,590	4,722	1,809	1,124	170,869	139,239	310,108	107.08	33.61
(%)	(2.0)	(9.4)	(20.8)	(24.8)	(20.0)	(12.3)	(6.2)	(2.8)	(1.1)	(0.7)	(100.0)				
記載なし															
(%)															
総計	3,410	16,101	35,569	42,378	34,203	20,963	10,590	4,722	1,809	1,124	170,869	139,239	310,108	107.08	33.61
(%)	(2.0)	(9.4)	(20.8)	(24.8)	(20.0)	(12.3)	(6.2)	(2.8)	(1.1)	(0.7)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査