

表 3759 Non-HDL-C濃度(mg/dL) 週透析回数(回/週)別 (透析患者全体)

週透析回数	50未満	50～	70～	90～	110～	130～	150～	170～	190～	210～	合計	記載なし	総計	平均	標準偏差
1回	10	59	145	257	255	177	99	51	15	14	1,082	758	1,840	117.68	34.72
(%)	(0.9)	(5.5)	(13.4)	(23.8)	(23.6)	(16.4)	(9.1)	(4.7)	(1.4)	(1.3)	(100.0)				
2回	77	286	727	832	660	460	253	112	54	30	3,491	2,701	6,192	109.25	35.43
(%)	(2.2)	(8.2)	(20.8)	(23.8)	(18.9)	(13.2)	(7.2)	(3.2)	(1.5)	(0.9)	(100.0)				
3回	3,260	15,405	33,897	40,118	32,074	19,416	9,704	4,282	1,608	978	160,742	109,897	270,639	106.55	33.32
(%)	(2.0)	(9.6)	(21.1)	(25.0)	(20.0)	(12.1)	(6.0)	(2.7)	(1.0)	(0.6)	(100.0)				
4回	15	53	84	157	145	95	52	21	11	6	639	355	994	113.32	35.15
(%)	(2.3)	(8.3)	(13.1)	(24.6)	(22.7)	(14.9)	(8.1)	(3.3)	(1.7)	(0.9)	(100.0)				
5回	1	1	6	7	17	12	4	2	3		53	36	89	124.94	32.97
(%)	(1.9)	(1.9)	(11.3)	(13.2)	(32.1)	(22.6)	(7.5)	(3.8)	(5.7)		(100.0)				
6回			4	11	10	4	3	1			33	48	81	116.79	24.60
(%)			(12.1)	(33.3)	(30.3)	(12.1)	(9.1)	(3.0)			(100.0)				
7回		1	3	5	7	3	1			2	22	8	30	125.23	46.36
(%)		(4.5)	(13.6)	(22.7)	(31.8)	(13.6)	(4.5)			(9.1)	(100.0)				
合計	3,363	15,805	34,866	41,387	33,168	20,167	10,116	4,469	1,691	1,030	166,062	113,803	279,865	106.71	33.40
(%)	(2.0)	(9.5)	(21.0)	(24.9)	(20.0)	(12.1)	(6.1)	(2.7)	(1.0)	(0.6)	(100.0)				
記載なし	47	296	703	991	1,035	796	474	253	118	94	4,807	25,436	30,243	119.84	38.06
(%)	(1.0)	(6.2)	(14.6)	(20.6)	(21.5)	(16.6)	(9.9)	(5.3)	(2.5)	(2.0)	(100.0)				
総計	3,410	16,101	35,569	42,378	34,203	20,963	10,590	4,722	1,809	1,124	170,869	139,239	310,108	107.08	33.61
(%)	(2.0)	(9.4)	(20.8)	(24.8)	(20.0)	(12.3)	(6.2)	(2.8)	(1.1)	(0.7)	(100.0)				
平均	2.98	2.98	2.97	2.97	2.97	2.97	2.96	2.96	2.96	2.96	2.97	2.97	2.97		
標準偏差	0.20	0.19	0.21	0.23	0.25	0.26	0.27	0.28	0.28	0.34	0.24	0.24	0.24		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査