

表 3766 Non-HDL-C濃度(mg/dL) 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	50未満	50～	70～	90～	110～	130～	150～	170～	190～	210～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	755	3,520	8,228	10,007	8,375	5,080	2,627	1,104	393	227	40,316	28,261	68,577	107.72	32.97
(%)	(1.9)	(8.7)	(20.4)	(24.8)	(20.8)	(12.6)	(6.5)	(2.7)	(1.0)	(0.6)	(100.0)				
慢性腎盂腎炎	23	105	217	284	262	162	77	43	10	2	1,185	980	2,165	108.92	32.38
(%)	(1.9)	(8.9)	(18.3)	(24.0)	(22.1)	(13.7)	(6.5)	(3.6)	(0.8)	(0.2)	(100.0)				
急速進行性糸球体腎炎	15	60	194	270	219	142	97	51	13	23	1,084	750	1,834	115.30	36.72
(%)	(1.4)	(5.5)	(17.9)	(24.9)	(20.2)	(13.1)	(8.9)	(4.7)	(1.2)	(2.1)	(100.0)				
妊娠腎/妊娠中毒症	3	27	95	153	146	101	57	24	11	7	624	440	1,064	117.56	33.38
(%)	(0.5)	(4.3)	(15.2)	(24.5)	(23.4)	(16.2)	(9.1)	(3.8)	(1.8)	(1.1)	(100.0)				
その他の分類不能の腎炎	10	59	127	149	105	66	33	17	4	5	575	374	949	106.13	34.44
(%)	(1.7)	(10.3)	(22.1)	(25.9)	(18.3)	(11.5)	(5.7)	(3.0)	(0.7)	(0.9)	(100.0)				
多発性嚢胞腎	74	367	990	1,240	1,108	682	299	145	28	27	4,960	3,238	8,198	108.67	31.77
(%)	(1.5)	(7.4)	(20.0)	(25.0)	(22.3)	(13.8)	(6.0)	(2.9)	(0.6)	(0.5)	(100.0)				
腎硬化症	198	1,120	2,715	3,504	2,675	1,526	712	258	135	73	12,916	8,544	21,460	106.41	31.76
(%)	(1.5)	(8.7)	(21.0)	(27.1)	(20.7)	(11.8)	(5.5)	(2.0)	(1.0)	(0.6)	(100.0)				
悪性高血圧	17	72	225	260	207	130	72	36	17	8	1,044	837	1,881	110.21	35.59
(%)	(1.6)	(6.9)	(21.6)	(24.9)	(19.8)	(12.5)	(6.9)	(3.4)	(1.6)	(0.8)	(100.0)				
糖尿病性腎症	1,270	5,923	12,022	13,228	9,865	5,877	2,936	1,365	551	353	53,390	37,117	90,507	104.41	33.79
(%)	(2.4)	(11.1)	(22.5)	(24.8)	(18.5)	(11.0)	(5.5)	(2.6)	(1.0)	(0.7)	(100.0)				
SLE腎炎	13	64	168	223	184	134	88	33	16	7	930	620	1,550	113.76	35.43
(%)	(1.4)	(6.9)	(18.1)	(24.0)	(19.8)	(14.4)	(9.5)	(3.5)	(1.7)	(0.8)	(100.0)				
アミロイド腎	11	29	41	39	34	20	11	5	4	1	195	130	325	100.97	38.40
(%)	(5.6)	(14.9)	(21.0)	(20.0)	(17.4)	(10.3)	(5.6)	(2.6)	(2.1)	(0.5)	(100.0)				
痛風腎	15	49	93	122	109	59	28	13	5	1	494	334	828	105.75	33.08
(%)	(3.0)	(9.9)	(18.8)	(24.7)	(22.1)	(11.9)	(5.7)	(2.6)	(1.0)	(0.2)	(100.0)				
先天性代謝異常による腎不全	5	10	19	18	18	10	8	3	2	2	95	82	177	108.01	40.58
(%)	(5.3)	(10.5)	(20.0)	(18.9)	(18.9)	(10.5)	(8.4)	(3.2)	(2.1)	(2.1)	(100.0)				
腎・尿路結核	1	9	21	15	28	8	7	2		1	92	66	158	106.73	32.50
(%)	(1.1)	(9.8)	(22.8)	(16.3)	(30.4)	(8.7)	(7.6)	(2.2)		(1.1)	(100.0)				
腎・尿路結石	4	32	49	67	51	31	16	6			256	174	430	103.86	29.79
(%)	(1.6)	(12.5)	(19.1)	(26.2)	(19.9)	(12.1)	(6.3)	(2.3)			(100.0)				
腎・尿路腫瘍	7	44	104	108	76	49	28	11	5	2	434	252	686	105.57	33.14
(%)	(1.6)	(10.1)	(24.0)	(24.9)	(17.5)	(11.3)	(6.5)	(2.5)	(1.2)	(0.5)	(100.0)				
閉塞性尿路障害	4	28	62	67	73	52	18	13	2	3	322	195	517	110.87	33.76
(%)	(1.2)	(8.7)	(19.3)	(20.8)	(22.7)	(16.1)	(5.6)	(4.0)	(0.6)	(0.9)	(100.0)				
骨髄腫		14	19	23	22	19	9	5	1	2	114	76	190	113.04	35.77
(%)		(12.3)	(16.7)	(20.2)	(19.3)	(16.7)	(7.9)	(4.4)	(0.9)	(1.8)	(100.0)				
腎形成不全	7	19	50	58	51	30	25	7	3		250	169	419	109.76	33.27
(%)	(2.8)	(7.6)	(20.0)	(23.2)	(20.4)	(12.0)	(10.0)	(2.8)	(1.2)		(100.0)				
不明	218	988	2,292	2,801	2,260	1,444	651	324	95	59	11,132	8,993	20,125	107.21	32.97
(%)	(2.0)	(8.9)	(20.6)	(25.2)	(20.3)	(13.0)	(5.8)	(2.9)	(0.9)	(0.5)	(100.0)				
再導入	13	81	161	229	166	124	60	25	6	8	873	544	1,417	108.93	34.08
(%)	(1.5)	(9.3)	(18.4)	(26.2)	(19.0)	(14.2)	(6.9)	(2.9)	(0.7)	(0.9)	(100.0)				
その他	53	286	640	714	645	387	184	77	37	21	3,044	2,470	5,514	107.75	33.65
(%)	(1.7)	(9.4)	(21.0)	(23.5)	(21.2)	(12.7)	(6.0)	(2.5)	(1.2)	(0.7)	(100.0)				
合計	2,716	12,906	28,532	33,579	26,679	16,133	8,043	3,567	1,338	832	134,325	94,646	228,971	106.43	33.32
(%)	(2.0)	(9.6)	(21.2)	(25.0)	(19.9)	(12.0)	(6.0)	(2.7)	(1.0)	(0.6)	(100.0)				
記載なし															
(%)															
総計	2,716	12,906	28,532	33,579	26,679	16,133	8,043	3,567	1,338	832	134,325	94,646	228,971	106.43	33.32
(%)	(2.0)	(9.6)	(21.2)	(25.0)	(19.9)	(12.0)	(6.0)	(2.7)	(1.0)	(0.6)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査